

CHAPTER 6

Chi-Square Test & Contingency Analysis

**(Chi-Square Test for k Proportions,
Chi-Square Test of Independence Contingency Table)**

Chi-Square Test &

One Way Contingency Table

Part 1: Chi-Square Analysis & How to find critical value

Part 2: Categories with Equal Frequencies/Probabilities

Part 3: Categories with Unequal Frequencies/Probabilities

Part 4: 2-Way Contingency Table

Multinomial Experiment

An experiment that meets the following conditions:

1. The number of trials is fixed.
2. The trials are independent.
3. All outcomes of each trial must be classified into exactly one of several different categories.
4. The probabilities for the different categories remain constant for each trial.

Multinomial Experiment (cont.)

- n identical trials *vs the experiments*
- k outcomes to each trial
- Constant outcome probability, p_k
- Independent trials
- Random variable is count, O_k

Multinomial Experiment (cont.)

Example:



shutterstock.com - 598513013

Sample (n)

100 people from Malaysia population



Categories / outcome (k)

9 candidates

Goodness-of-fit Test

Goodness-of-fit test is used to test the hypothesis that an observed frequency distribution fits (or conforms to) some claimed distribution.⁽²⁾

*expected distribution /
expected frequency for
the test*

Goodness-of-fit Test (cont.)

Notation:

- O represents the observed frequency of an outcome
- E represents the expected frequency of an outcome
- k represents the number of different categories or outcomes
- n represents the total number of trials

Expected Frequencies

If all expected frequencies are **equal**:

$$E = \frac{n}{k}$$

the sum of all observed frequencies divided by the number of categories.

Expected Frequencies (cont.)

If all expected frequencies are **not all equal**:

$$E = n * p$$

each expected frequency is found by **multiplying** the sum of all observed frequencies (n) by the probability for the category (p).

Expected Frequencies (cont.)

Key Question : *of goodness of fit test*

Are the differences between the **observed values (O)** and the theoretically **expected values (E)** statistically significant?

Answer:

We need to measure the discrepancy between **O** and **E**; the test statistic will involve their difference: **O - E**

Chi-Square Test


$$\chi^2 = \sum \frac{(O - E)^2}{E}$$



Test statistic value->
calculated.

Critical Values (Chi-square value from table):

1. Found in table χ^2 using $k-1$ degrees of freedom where k = number of categories.
2. Goodness-of-fit hypothesis tests are always right-tailed.



Test Hypothesis

$$H_0 : \mu_1 = \mu_2$$
$$H_1 : \mu_1 \neq \mu_2$$

H_0 : No difference between observed and expected probabilities.

H_1 : At least one of the probabilities is different from the others.

- A **close agreement** between observed and expected values will lead to a small value of χ^2 and a large p -value.
- A **large disagreement** between observed and expected values will lead to a large value of χ^2 and a small p -value.
- A significantly large value of χ^2 will cause a rejection of the null hypothesis of no difference between the observed and the expected.



Swiftness of fate

Relationships Among Components in Goodness-of-Fit Hypothesis Test

Compare the observed O values to the corresponding expected E values.

*O*s and *E*s are close.

*O*s and *E*s are far apart.



Fail to reject H_0

Reject H_0

Good fit with assumed distribution

Not a good fit with assumed distribution

Chi-Square (χ^2) Test for k Proportions

- Tests Equality (=) of Proportions Only
 - Example: $p_1 = 0.2$, $p_2 = 0.3$, $p_3 = 0.5$
- One variable with several levels.
- Assumptions:
 - Multinomial Experiment
 - Large Sample Size
 - **All expected counts ≥ 5**
- Uses One-Way Contingency Table

One-Way Contingency Table

- Shows number of observations in k Independent Groups (Outcomes or Variable Levels)

Outcomes ($k = 3$)

1st row

Outcome →

Candidate			
Tom	Bill	Mary	Total
35	20	45	100

2nd row

response →

Number of responses


n, number of trials.

Finding Critical Value

Example: What is the critical χ^2 value if $k = 3$, and $\alpha = 0.05$?

If $o_i = e_i$, $\chi^2 = 0$. Do not reject H_0

$Df = k - 1 = 2$
 $Df_{2, 0.05} =$



χ^2 Table (Portion)	Upper Tail Area				
DF	.9959505
1	0.004	...	3.841
2	0.010	...	0.103	...	5.991

$Df = k - 1 = 2$ →

Categories With Equal Frequencies/Probabilities (Part 2)

Categories with Equal Frequencies/Probabilities


Statement of test hypothesis:

① $H_0: p_1 = p_2 = p_3 = \dots = p_k$ ← no diff

$H_1:$ at least one of the probabilities is different from the others.

Example 1

A study was conducted on ⁿ147 cases of industrial accidents that required medical attention. Test the claim that the accidents occur with equal proportions on the 5 workdays. (k)

Day	Mon	Tues	Wed	Thurs	Fri	$\rightarrow (k)$
Observed accidents	31	42	18	25	31	$\rightarrow 0$
						

Example 1 - Solution

Claim: Accidents occur with the same proportion. Therefore,

$$p_1 = p_2 = p_3 = p_4 = p_5$$

① Construct hypotheses

i. State the test hypothesis:

H_0 : $p_1 = p_2 = p_3 = p_4 = p_5$ ✓ = no diff

H_1 : At least 1 of the 5 proportions is different from others.

Example 1 – Solution (cont.)

ii. Calculate the expected frequency:

$$E = n/k = 147/5 = \underline{29.4}$$

$$E = \frac{n}{k}$$

- ① hypothesis ✓
- ② Expect ✓
- ③ Chi Sq ✓
- ④ critical value ✓
- ⑤ sign ✓
- ⑥ conclude

Observed and Expected Frequencies

Day	Mon	Tues	Wed	Thurs	Fri
O: Observed accidents	31	42	18	25	31
E: Expected accidents	29.4	29.4	29.4	29.4	29.4

Example 1 – Solution (cont.)

iii. Calculate the different between O and E:

$$\chi^2 = (O - E)^2 / E$$

Observed and Expected Frequencies of Industrial Accidents

Day	Mon	Tues	Wed	Thurs	Fri
Observed accidents	31	42	18	25	31
Expected accidents	29.4	29.4	29.4	29.4	29.4
$(O - E)^2 / E$	<u>0.0871</u>	<u>5.4000</u>	<u>4.4204</u>	0.6585	0.0871 (rounded)

$$\frac{(O - E)^2}{E} = \frac{(31 - 29.4)^2}{29.4} = 0.0871$$

Example 1 – Solution (cont.)

Observed and Expected Frequencies of Industrial Accidents

Day	Mon	Tues	Wed	Thurs	Fri
Observed accidents	31	42	18	25	31
Expected accidents	29.4	29.4	29.4	29.4	29.4
$(O - E)^2/E$	0.0871	5.4000	4.4204	0.6585	0.0871 (rounded)


iv. Calculate the test statistic: (calculated chi-square value)

$$\chi^2 = \sum \frac{(O - E)^2}{E} = 0.0871 + 5.4000 + 4.4204 + 0.6585 + 0.0871 = \underline{10.6531} \quad \checkmark$$


Example 1 – Solution (cont.)

v. Find the critical value: (chi-square value from table)

- Refer to table χ^2 with $k-1 = 5 - 1 = 4$; and $\alpha = 0.05$.
- It will show that $\chi^2_{4,0.05} = 9.48$



Example 1 – Solution (cont.)



vi. Based on the result, state the conclusion:

Test statistic falls within the critical region, therefore we reject hypothesis null. That is, we reject claim that the accidents occur with equal proportions (frequency) on the 5 workdays.

Example 2

As personnel director, you want to test the perception of fairness of three methods of performance evaluation.

Of 180ⁿ employees,

- o **63** rated **Method 1** as fair.
 - o **45** rated **Method 2** as fair.
 - o **72** rated **Method 3** as fair.
- } K

- ① hypothesis
- ② Expect for
- ③ Chi square
- ④ Control case
- ⑤ group
- ⑥ course

At the 0.05 level, is there a difference in perceptions?

$\alpha = 0.05$



Example 2 – Solution (cont.)


i. Test hypothesis:

$$H_0: p_1 = p_2 = p_3 = 1/3$$

H_1 : At least 1 is different

ii. Find the critical value:

$$\alpha = 0.05; k = 3 - 1 = 2$$




Example 2 – Solution (cont.)

- iii. Calculate the expected counts and,
- iv. Find the test statistics value:

Cell, i	Observed Count, o_i	Expected Count, e_i	$[o_i - e_i]^2 / e_i$
1	63	$(1/3) \times 180 = 60$	0.15
2	45	$(1/3) \times 180 = 60$	3.75
3	72	$(1/3) \times 180 = 60$	2.40
Total	180	180	$\chi^2 = 6.30$

Example 2 – Solution (cont.)

v. State the decision:



Test Statistic: $\chi^2 = 6.3$

Critical value: $\chi^2 (k=2, \alpha=0.05)$
= 5.991

Conclusion:

Reject H_0 at $\alpha = .05$

There is evidence of a difference in proportions.

Example 3

- Are technical support calls equal across all days of the week?

■ Sample data:

	<u>Sum of calls for each day:</u>
Monday	290
Tuesday	250
Wednesday	238
Thursday	257
Friday	265
Saturday	230
Sunday	192
	$\Sigma = 1722$

Example 3 – Solution

- If calls are equal across all days of the week, the 1722 calls would be expected to be equally divided across the 7 days:

$$E = \frac{n}{k}$$

$$E = \frac{1722}{7} = 246 \text{ expected calls per day}$$

i. Test hypothesis:

$$H_0: p_1 = p_2 = p_3 = p_4 = p_5 = p_6 = p_7 = 1/7 \checkmark$$

H_1 : At least 1 is different \checkmark

Example 3 – Solution (cont.)

- ii. Calculate the expected counts and,
 iii. Find the test statistics value:

②
 ③

	Observed o_i	Expected e_i	$\chi^2 = [o_i - e_i]^2 / e_i$
Monday	290	246	7.8699
Tuesday	250	246	0.0650
Wednesday	238	246	0.2602
Thursday	257	246	0.4919
Friday	265	246	1.4675
Saturday	230	246	1.0407
Sunday	192	246	11.8537
TOTAL	1722	1722	$\chi^2 = 23.0489$

Example 3 – Solution (cont.)

iv. Find the critical value: $k = 7$


- $k - 1 = \underline{6}$ (7 days of the week) so use 6 degrees of freedom

$$\chi^2_{.05,6} = \underline{12.592}$$

v. State the decision: (6)

Conclusion:

$\chi^2 = 23.0489 > \chi^2_{\alpha} = 12.592$ so **reject H_0** and conclude that the distribution is not uniform.



Categories With Unequal Frequencies/Probabilities (Part 3)

Categories with **Unequal** Frequencies/Probabilities

H_0 : $p_1, p_2, p_3, \dots, p_k$ are as claimed.

H_1 : at least one of the above proportions is
different from the claimed value.

Example 4

Mars, Inc. claims its M&M candies are distributed with the color percentages of 30% brown, 20% yellow, 20% red, 10% orange, 10% green, and 10% blue. At the 0.05 significance level, test the claim that the color distribution is as claimed by Mars, Inc. The observed frequency as shown below:

Frequencies of M&Ms candies

	Brown	Yellow	Red	Orange	Green	Blue	$\rightarrow k = 6$
Observed frequency	33	26	21	8	7	5	$\rightarrow 0$

Example 4 - Solution

Claim: $p_{\text{brown}} = 0.30$, $p_{\text{yellow}} = 0.20$, $p_{\text{red}} = 0.20$,
 $p_{\text{orange}} = 0.10$, $p_{\text{green}} = 0.10$, $p_{\text{blue}} = 0.10$

i. **Statement of test hypothesis:**

H_0 : $p_{\text{brown}} = 0.30$, $p_{\text{yellow}} = 0.20$, $p_{\text{red}} = 0.20$,
 $p_{\text{orange}} = 0.10$, $p_{\text{green}} = 0.10$, $p_{\text{blue}} = 0.10$.

H_1 : At least one of the proportions is different from the claimed value.

ii. Calculate the expected frequency:

2

Frequencies of M&Ms candies

	Brown	Yellow	Red	Orange	Green	Blue
Observed frequency	33	26	21	8	7	5

$n = \underline{\underline{100}}$

Expected frequency:

Brown $E = np = (100)(0.30) = 30$

Yellow $E = np = (100)(0.20) = 20$

Red $E = np = (100)(0.20) = 20$

Orange $E = np = (100)(0.10) = 10$

Green $E = np = (100)(0.10) = 10$

Blue $E = np = (100)(0.10) = 10$

$E = n * p$

3

iii. Calculate the test statistic @chi-square value:

	Brown	Yellow	Red	Orange	Green	Blue	
Observed frequency	33	26	21	8	7	5	✓
Expected frequency	30	20	20	10	10	10	✓
$\chi^2 (O - E)^2 / E$	0.3	1.8	0.05	0.4	0.9	2.5	

Test statistics value:

$$\chi^2 = \sum \frac{(O - E)^2}{E} = \underline{\underline{5.95}}$$

iv. Find the critical value from chi-square table:


Critical Value $\chi^2 = \underline{11.071}$
(with $k-1 = 5$ and $\alpha = 0.05$)

4

v. State the decision:

Test statistic value ($\chi^2 = 5.95$) < critical value ($\chi^2_{k=5, \alpha=0.05} = 11.071$),
that is it does not fall within critical region. Thus, we do not reject H_0 .

5



There is not sufficient evidence to warrant rejection of the claim that the colors are distributed with the given percentages.

Exercise

It was claimed that population at ABC country in 2008 consisted of 50.7% English, 6.6% French, 30.6% Irish, 10.8% Asians, and 1.3% other ethnic groups. Suppose that a random sample of 1000 student graduating from ABC colleges and universities in 2008 resulted in the accompanying data on ethnic group (see table below).

Ethnic Group	Number in Sample
English	679
French	51
Irish	77
Asian	190
Other	3

Do the data provide evidence that the proportion of students graduating from colleges and universities in ABC for these ethnic group categories differs from the respective proportions in the population for ABC? Test the appropriate hypotheses using $\alpha=0.01$.

Chi-Square Test & Two Way Contingency Table (Part 4)

Chi-Square (χ^2) Test of Independence

- To show if a relationship exists between 2 qualitative variables, when
 - One sample is drawn.
 - Does **not** show causality.
- Assumptions:
 - Multinomial experiment.
 - All expected counts ≥ 5
- Uses two-way contingency table

Recap: One-Way Contingency Table

- Shows number of observations in k Independent Groups (Outcomes or Variable Levels)

Outcomes ($k = 3$)

		Candidate			
k		Tom	Bill	Mary	Total
O		35	20	45	100

Number of responses

Two-Way Contingency Table

- Shows # observations from 1 sample jointly in 2 qualitative variables:

House Style	House Location		Total
	Urban	Rural	
Split-Level	63	49	112
Ranch	15	33	48
Total	78	82	160

Level of variable: 2 (points to House Location)

Level of variable: 1 (points to House Style)


Test hypotheses & Test Statistic

- Test hypothesis:

H_0 : Variables are independent.

H_1 : Variables are related (dependent).

- Test Statistic:

$$\chi^2 = \sum_{\text{all cells}} \frac{\left[\overset{\text{Observed count}}{o_{ij}} - \underset{\text{Expected count}}{e_{ij}} \right]^2}{e_{ij}}$$


- Degrees of Freedom: $(r - 1)(c - 1)$

df.

Calculation of Expected Counts

- Statistical independence means joint probability equals product of marginal probabilities.
- Compute marginal probabilities & multiply for joint probability.
- Expected count is sample size times joint probability.

Example 5: How to Calculate Expected Counts

② House Style	① Location		Total
	Urban Obs.	Rural Obs.	
Split-Level	63	49	112 ✓
Ranch	15	33	48 ✓
Total	78	82	160

Handwritten annotations: A red box highlights the 63 and 49 cells with a '+' sign between them. A red arrow points from this box to the 112 total. Another red box highlights the 33 cell with a '+' sign above it, and a red arrow points from this box to the 82 total. A red double underline is under the 82 total.

Example 5 (cont.)

Marginal probability = $\frac{112}{160}$ *total row*
n

House Style	Location		Total
	Urban Obs.	Rural Obs.	
Split-Level	63	49	112
Ranch	15	33	48
Total	<u>78</u>	82	<u>160</u>

Handwritten notes:

- Blue arrow points from the 'Total' column header to the '112' value in the 'Split-Level' row.
- Red arrow points from the '112' value to the fraction $\frac{112}{160}$.
- Red arrow points from the '48' value to the fraction $\frac{48}{160}$.
- Red arrow points from the '78' value to the fraction $\frac{78}{160}$.
- Red arrow points from the '82' value to the fraction $\frac{82}{160}$.
- The '160' value in the 'Total' row is circled in red.

Example 5 (cont.)

Marginal probability = $\frac{112}{160}$

House Style	Location		Total
	Urban Obs.	Rural Obs.	
Split-Level	63	49	112
Ranch	15	33	48
Total	78	82	160

Marginal probability = $\frac{78}{160}$

Example 5 (cont.)

$$\text{Joint probability} = \frac{112}{160} \times \frac{78}{160}$$

$$\text{Marginal probability} = \frac{112}{160}$$

House Style	Location		Total
	Urban Obs.	Rural Obs.	
Split-Level	63	49	112
Ranch	15	33	48
Total	78	82	160

$$\text{Marginal probability} = \frac{78}{160}$$

Example 5 (cont.)

$$\begin{aligned} \text{Expected } E &= \frac{h}{k} \\ \text{Unex} &= n \cdot p \end{aligned}$$

Expected Count calculation formula:

$$e_{ij} = \frac{(i^{\text{th}} \text{ Row total})(j^{\text{th}} \text{ Column total})}{\text{Total sample size}}$$

Example 5 (cont.)

$$\text{Joint probability} = \frac{112}{160} \times \frac{78}{160}$$

$$\text{Marginal probability} = \frac{112}{160}$$

House Style	Location		Total
	Urban Obs.	Rural Obs.	
Split-Level	63	49	112
Ranch	15	33	48
Total	78	82	160

$$\text{Marginal probability} = \frac{78}{160}$$

Example 5 (cont.)

House Style	House Location				Total
	Urban	Rural	Urban	Rural	
	Obs.	Exp.	Obs.	Exp.	
Split-Level	63	<u>54.6</u>	49	<u>57.4</u>	112
Ranch	15	<u>23.4</u>	33	<u>24.6</u>	48
Total	78	78	82	82	160

$\frac{112 \cdot 78}{160}$ $\frac{112 \cdot 82}{160}$
 $\frac{48 \cdot 78}{160}$ $\frac{48 \cdot 82}{160}$

Example 6

n You're a marketing research analyst. You ask a random sample of 286 consumers if they purchase Diet Pepsi or Diet Coke. At the 0.05 α level, is there evidence of a relationship? V_1 V_2

Diet Coke	Diet Pepsi		Total
	No	Yes	
No	84	32	116
Yes	48	122	170
Total	132	154	286

C_1

C_2

R_1
 R_2
 R_3

Example 6 - Solution

①

i. State the test hypothesis:

H_0 : No relationship between variables.


H_1 : Variables has relationship.

ii. Find the critical value (refer to chi-square table):

$\alpha = 0.05$

$df = (2 - 1)(2 - 1) = 1$

④ $\chi^2_{0.05, 1} = 3.841$



Example 6 – Solution (cont.)

iii. Calculate the expected counts: 2

		Diet Pepsi				
		No		Yes		
Diet Coke	Obs.	Exp.	Obs.	Exp.	Total	
No	84	53.5	32	62.5	116	$\frac{116 \cdot 132}{286}$
Yes	48	78.5	122	91.5	170	$\frac{170 \cdot 154}{286}$
Total	132	132	154	154	286	$\frac{170 \cdot 132}{286}$

✓ $e_{ij} \geq 5$ in all cells

$$\text{Expected} = \frac{(\text{Row total}) \times (\text{Col total})}{n}$$

Example 6 – Solution (cont.)


iv. Calculate the test statistic value:

Cell, ij	Observed Count, o_{ij}	Expected Count, e_{ij}	$\chi^2 = [o_{ij} - e_{ij}]^2 / e_{ij}$
1,1	84	$(116)(132)/286$ $=53.5$ ✓	17.39
1,2	32	$(116)(154)/286$ $=62.5$ ✓	14.88
2,1	48	$(170)(132)/286$ $=78.5$ ✓	11.85
2,2	122	$(170)(154)/286$ $=91.5$ ✓	10.17
			$\chi^2 =$ 54.29

Example 6 – Solution (cont.)

v. State the decision:

⑤ graph



Test Statistic: $\chi^2 = 54.29$

Critical value: $\chi^2_{k=1, \alpha=0.05} = 3.841$

Decision:

Since, test statistic value $>$ critical value, thus reject H_0 at $\alpha = 0.05$

Conclusion:

There is evidence of a relationship between the variables.

Example 7

- Left-Handed vs. Gender
 - Dominant Hand: Left vs. Right
 - Gender: Male vs. Female

H_0 : Hand preference is independent of gender

H_1 : Hand preference is **not** independent of gender

Example 7 – Solution

- Sample results organized in a contingency table:

sample size = $n = 300$:

120 Females, 12 were left handed

180 Males, 24 were left handed

Gender	Hand Preference		
	Left	Right	
Female	12	108	120
Male	24	156	180
	36	264	300

Example 7 – Solution (cont.)

- Observed frequencies vs. expected frequencies:

Gender	Hand Preference		
	Left	Right	
Female	Observed = 12 Expected = 14.4	Observed = 108 Expected = 105.6	120
Male	Observed = 24 Expected = 21.6	Observed = 156 Expected = 158.4	180
	36	264	300

Example 7 – Solution (cont.)

Cell, ij	Observed Count, o_{ij}	Expected Count, e_{ij}	$[o_{ij} - e_{ij}]^2 / e_{ij}$
1,1	12	$(120)(36)/300$ =14.4	0.4000
1,2	108	$(120)(264)/300$ =105.6	0.0545
2,1	24	$(180)(36)/300$ =21.6	0.2667
2,2	156	$(180)(264)/300$ =158.4	0.0364
$\chi^2 =$			0.7576

Example 7 – Solution (cont.)

Test Statistic: $\chi^2 = 0.7576$


Critical value: $\chi^2_{k=1, \alpha = 0.05} = 3.841$

Decision:

Since, test statistic value $<$ critical value, thus do not reject H_0 at $\alpha = 0.05$

Conclusion:

There is evidence that gender and hand preference are independent.



Solution

Cell, ij	Observed Count, o_{ij}	Expected Count, e_{ij}	$[o_{ij} - e_{ij}]^2 / e_{ij}$
1,1	117	91.5	7.11
1,2	66	91.5	7.11
2,1	150	155	0.16
2,2	160	155	0.16
3,1	109	138.5	6.28
3,2	168	138.5	6.28
4,1	124	115	0.70
4,2	106	115	0.70
$\chi^2 =$			28.51

Example 7 – Solution (cont.)

Test Statistic: $\chi^2 = 28.51$

Critical value: $\chi^2_{k=3, \alpha = 0.05} = 7.815$

Decision:

Since, test statistic value $>$ critical value, thus we reject H_0 at $\alpha = 0.05$

Conclusion:

There is evidence that male and female inmates differ with respect to type of offense.

Exercise

Jail inmates can be classified into one of the following four categories according to the type of crime committed: violent crime, crime against property, drug offenses, and public-order offenses. Suppose that random samples of 500 male inmates and 500 female inmates are selected, and each inmate is classified according to type of offense.

Type of Crime	Gender	
	Male	Female
Violent	117	66
Property	150	160
Drug	109	168
Public-order	124	106

We would like to know whether male and female inmates differ with respect to type of offense. Test the relevant hypotheses using a significance level of 0.05.