

SECI2143: PROBABILITY & STATISTICAL DATA ANALYSIS

CHAPTER 2

Data Description



Types of Data Set

Univariate

- Multivariate
 - Bivariate



Univariate Data Set

- Univariate data set consists of observations on a single variable made on individuals in a sample or population.
- **Variable** is any characteristic whose value may change from one individual or object to another.
- A univariate data set is **categorical** (or qualitative) if the individual observations are categorical response.
 - Example: recorded calculator brand.
- A univariate data set is numerical (or quantitative) if each observation is a number.
 - Example: recorded number of calculator purchased.



Multivariate Data Set

- Multivariate data set consists of observations on two or more variables made on individuals in a sample or population.
 - Example: recorded height, weight, pulse rate and systolic blood pressure for each individual in a group.
- A bivariate data are special case of multivariate data set because it focus simultaneously on only two variables or two different characteristics.
 - Example: recorded both height and weight for each individual in a group.



Categorical Data

(when numbers are collected in groups or categories)



Displaying Categorical Data

- An appropriate graphical or tabular of data can be an effective way to summarize and communicate information.
- Example:
 - Frequency distribution or frequency table
 - Bar chart
 - Pie chart
 - Dot-Plot



Frequency Distribution

- A frequency distribution for categorical data is a table that displays the possible categories along with the associated frequencies and/or relative frequencies.
- The frequency for a particular category is the number of times the category appears in the data set.
- The **relative frequency** for a particular category is the fraction or proportion of the observations resulting in the category.

Relative frequency=
$$\frac{\text{frequency}}{\text{Total no. of cases}}$$

 Frequency distributions are portrayed as frequency tables, dot plot, histograms, or polygons.



Table 1: Data of staff at faculty of computing

Staff	Position	Blood Type	Weight	Height	Qualification
1	Senior Lecturer	Α	60	165	PhD
2	Lecturer	В	55	150	Master
3	Professor	0	65	170	PhD
4	Associate Professor	AB	70	175	PhD
5	Associate Professor	0	61	160	PhD
6	Senior Lecturer	0	58	155	PhD
7	Senior Lecturer	В	48	167	PhD
8	Lecturer	Α	68	174	Master
9	Lecturer	Α	55	150	Master
10	Associate Professor	AB	62	163	PhD
11	Professor	0	58	165	PhD
•••		•••	•••	•••	
140	Tutor	0	45	150	Master





Based on data in Table 1, we can summarize the info using the **Frequency Table.**

Table 2: Staff distribution in faculty of computing based on position

Position	Number of Staff
	(frequency)
Professor	12
Associate Professor	20
Senior Lecturer	59
Lecturer	40
Tutor	9
Total	140





Next, based on frequency table, we can calculate the **relative frequency**: (the ratio of the frequency of a particular event in a statistical experiment to the total frequency)

Position	frequency	Relative frequency
Professor	12	12 ÷ 140 = 0.09
Associate Professor	20	0.14
Senior Lecturer	59	0.42
Lecturer	40	0.29
Tutor	9	0.06
Total	140	1.00



Raw Data – A survey on the Internet usage by age

Age	NetUse	Age	NetUse	Age	NetUse
26.0	Yes	45.0	No	55.0	No
48.0	Yes	19.0	No	37.0	No
67.0	Yes	82.0	No	43.0	Yes
44.0	No	83.0	No	29.0	Yes
52.0	No	20.0	Yes	57.0	Yes
52.0	No	89.0	No	36.0	No
51.0	Yes	88.0	Yes	52.0	No
52.0	No	72.0	Yes	56.0	Yes
77.0	No	82.0	Yes	66.0	Yes
40.0	No	34.0	No	46.0	No



Example (cont'd)

List the variables involved and calculate the frequency:

Response	Frequency
No	17
Yes	13

- There could be another types of response with missing value. For example:
 - DK (Don't know)
 - NAP (Not applicable)
 - NA (No answer/response)



Example (cont'd)

Calculate the percent, valid percent and cumulative percent:

Response	Frequency	Percent	Valid Percent	Cumulative Percent
No	17	56.7	56.7	56.7
Yes	13	43.3	43.3	100.0
Total	30	100.0	100.0	

 Valid percent: excludes data with missing value (i.e., DK, NAP, and NA)



Exercise #1

The data below represent the color of M&Ms chocolate in a bag. Construct a relative frequency distribution.

Yellow	Orange	Brown	Green	Green
Blue	Brown	Red	Brown	Brown
Orange	Brown	Red	Brown	Red
Green	Brown	Red	Green	Yellow
Yellow	Red	Red	Brown	Orange
Yellow	Orange	Red	Orange	Blue
Brown	Red	Yellow	Brown	Red
Brown	Yellow	Yellow	Blue	Yellow
Yellow	Brown	Yellow	Green	Orange




Bar Chart


- A bar chart is a graph of the frequency distribution of categorical data.
- Each category in the frequency distribution is represented by a bar or rectangle.
- How to construct:
 - Draw a horizontal line, and write the category names or labels below the line at regularly spaced intervals.
 - Draw a vertical line, and label the scale using either frequency or relative frequency.
 - Place a rectangular bar above each category label. The height is determined by the category's frequency or relative frequency, and all bars should have the same width.
- What to look for:
 - Frequently and infrequently categories.













Staff Position



Example (cont'd)










Exercise #1 - SOLUTION

Color Distribution in M&Ms chocolate





Comparative Bar Charts


- Bar chart can also be used to give a visual comparison of two or more groups.
- When constructing a comparative bar graph we use the relative frequency rather than the frequency to construct the scale on the vertical axis so that we can make meaningful comparisons even if the sample sizes are not the same.



	Freq	luency	Relative Frequency		
Source of funding	Female	Male	Female	Male	
Student Loans	50	30	0.5	0.50	
Parents and/or family	30	20	0.3	0.33	
Scholarships	20	10	0.2	0.17	
Total	100	60	1.0	1.0	




Frequency



Inaccurate comparative bar chart

Relative frequency



Accurate comparative bar chart



Pie Charts

- In a pie chart, a circle is used to represent the whole data set, with "slices" of the pie representing the possible categories.
- The size of the slice for a particular category is proportional to the corresponding frequency or relative frequency.
- Pie charts are most effective for summarizing data sets when there are not too many different categories.




- How to construct
 - Draw a circle to represent the entire data set
 - For each category, calculate the "slice" size.
 slice size = 360 (category relative frequency)
 - Draw a slice of appropriate size for each category.


- What to look for
 - Categories that form large and small proportions of the data set.



Number of Students for Year 2013/2014 Intake




Percentage of Number of Students for Year 2013/2014 Intake





MAIN REASON FOR MIGRATION TO/FROM THE UK - 2007



SOURCE: ONS



Exercise #2

Travel Survey

92 people were asked how they got to work. Summarize of the data as follows:

- 35 used a car
- 42 took public transport
- 8 rode a bicycle
- 7 walked
- i) Calculate the relative frequency.
- ii) Draw the appropriate chart.



Numerical Data

(observations that can be measured)



Displaying Numerical Data

- Ungrouped data comprise a listing of the observed values.
- Grouped data represent a lumping together of the observed values.
- The data can be discrete or continuous.



Ungrouped data (discrete/integer):

```
      0
      1
      3
      0
      1
      0
      1
      0

      1
      5
      4
      1
      2
      1
      2
      0

      1
      0
      2
      0
      0
      2
      0

      2
      1
      1
      .....
```



Frequency distribution based on the grouped data:

Data	Frequency
0	15
1	20
2	8
3	5
:	:
:	:



Ungrouped data (continuous/fixed decimal):

2.559	2.556	2.566	2.546	2.561	2.570	2.546
2.565	2.543	2.538	2.560	2.560	2.545	2.551
2.568	2.546	2.555	2.551	2.554	2.574	2.568
2.572	2.550	2.556	2.551	2.561	2.560	2.564
2.567	2.560	2.551	2.562	2.542	2.549	2.561



Frequency distribution based on the grouped data:

Data	Frequency
2.531 - 2.535	6
2.536 - 2.540	8
2.541 – 2.545	12
2.546 – 2.550	13
:	:
:	:



Frequency Distributions

- A frequency distribution shows the number of observations falling into each of several ranges of values.
- Frequency distributions are portrayed as frequency tables, histograms, or polygons.
- Frequency distributions can show either the actual number of observations falling in each range or the percentage of observations. In the later instance, the distribution is called a relative frequency distribution.



• Discrete Data:

5	7	7	1
3	2	8	6
8	2	4	4
9	10	2	6
3	1	6	6
9	9	7	5
7	10	8	1
5	8		

Marks	Tally	Frequency
1	///	3
2	///	3
3	//	2
4	//	2
5	//	2
6	HHT .	5
7	////	4
8	HHT .	5
9	//	2
10	//	2
Total		30



Frequency distribution table (details):

Marks	Frequency	Relative frequency	Cumulative frequency	Percent(%) of CF
			(CF)	
1	3	0.100	0.1	10.0
2	3	0.100	0.2	20.0
3	2	0.067	0.267	26.7
4	2	0.067	0.334	33.4
5	3	0.100	0.434	43.4
6	4	0.133	0.567	56.7
7	4	0.133	0.7	70.0
8	4	0.133	0.833	83.3
9	3	0.100	0.933	93.3
10	2	0.067	1.00	100.0
Total	30	1.0	100.0	



Frequency distribution table for grouped data (discrete data):

Frequency Distribution Tables - Grouped Data

Arrange the following data into a frequency distribution table:

65 73 64 85 66 77 82 93 86 63 58 63 62 79 61 74

Class	Class centre CC	Tally	Frequency <i>f</i>
50-59	54.5	I	1
60-69	64.5	1111	7
70-79	74.5	1111	4
80-89	84.5		3
90-99	94.5	1	1

Add up frequency column ≠16

Count scores

Double check



Continuous data:

5.1 14.6 10.3 17.0

12.1 18.3 21.4 15.4

22.4 29.7 15.3 19.5

23.3 28.1 16.9 24.9

Class Interval	Frequency
5 to <10	1
10 to <15	3
15 to <20	6
20 to <25	4
25 to <30	2



• Frequency table for grouped (continuous data):

HEIGHT OF PUPILS (CM)	FREQUENCY		
150 ≤ x < 55	2		
155 ≤ x < 160	6		
160 ≤ x < 165	9		
165 ≤ x < 170	5		
170 ≤ x < 175	1		



Dotplots

When to use: Small numerical data sets.

How to construct:

- Draw a horizontal line and mark it with an appropriate measurement scale.
- 2. Locate each value in the data set along the measurement scale, and represent it by a dot. If there are two or more observations with the same value, stack the dots vertically.

What to look for:

- ✓ A representative or typical value in the data set.
- ✓ The extent to which the data values spread out.
- ✓ The nature of the distribution of values along the number line.
- ✓ The presence of unusual values in the data set.




- A graphical data using dots.
- **Example**: Minutes To Eat Breakfast

A survey of "How long does it take you to eat breakfast?" has these results:

Minutes: **0 1 2 3 4 5 6 7 8 9 10 11 12** People: 6 2 3 5 2 5 0 0 2 3 7 4 1

Which means that 6 people take 0 minutes to eat breakfast (they probably had no breakfast!), 2 people say they only spend 1 minute having breakfast, etc. The dot plot shown below,





Histograms

- A histogram is the most commonly used graph to show frequency distributions.
- A histogram consists of a set of rectangles that represent the frequency in each categories.
- It represents graphically the frequencies of the observed values.
- What to look for:
 - Central or typical value
 - Extent of spread or variation
 - General shape
 - Location and number of peaks
 - Presents of gap and outliers



How to construct:

- 1 Collect data and construct a tally sheet:
 - The number of cells should be between 5 and 20.
 - Use 5 to 9 cells when the number of observation <100.
 - Use 8 to 17 (between 100 and 500).
 - Use 15 to 20 (>500).
- 2 Determine the range:
 - $\blacksquare R = X_h X_l$
 - Where R = range, X_h = highest number, X_l = lowest number



- 3 Determine the cell interval:
 - The distance between adjacent cell midpoints.
 - An odd interval is recommended, so that the midpoint values will be the same number of decimal places as the data values.
 - Sturgis' rule:

$$i = \frac{R}{1 + 3.322 \log n}$$

- (n = number of observations)
- Trial and error, h=R/i
 (h=number of cells, R=range)



- 4 Determine the cell midpoint:
 - The lowest cell midpoint must be located to include the lowest data value in its cell.
 - The simplest technique is to select the lowest data point as the midpoint value for the first cell.
 - Use formula:

$$MP_l = X_l + \frac{i}{2}$$


(do not round-up/down the answer)

 MP_l = midpoint for lowest cell



- 5 Determine the cell boundaries:
 - Cell boundaries are the extreme or limit values of a cell (upper boundary and lower boundary)
 - All the observations that fall between the upper and lower boundaries are classified into that particular cell.
 - The boundary values are an extra decimal place or significant figure in accuracy than the observed values
- 6 Post the cell frequency.
- 7 Construct the histogram.







Example: Data of Steel Shaft Weight (kg)

Draw the histogram for the following data.

2.559	2.556	2.566	2.546	2.561	2.570	2.546
2.565	2.543	2.538	2.560	2.560	2.545	2.551
2.568	2.546	2.555	2.551	2.554	2.574	2.568
2.572	2.550	2.556	2.551	2.561	2.560	2.564
2.567	2.560	2.551	2.562	2.542	2.549	2.561
2.556	2.550	2.561	2.558	2.556	2.559	2.557
2.532	2.575	2.551	2.550	2.559	2.565	2.552
2.560	2.534	2.547	2.569	2.559	2.549	2.544
2.550	2.552	2.536	2.570	2.564	2.553	2.558
2.538	2.564	2.552	2.543	2.562	2.571	2.553
2.539	2.569	2.552	2.536	2.537	2.532	2.552
2.575 (highest)	2.545	2.551	2.547	2.537	2.547	2.533
2.538	2.571	2.545	2.545	2.556	2.543	2.551
2.569	2.559	2.534	2.561	2.567	2.572	2.558
2.542	2.574	2.570	2.542	2.552	2.551	2.553
2.546	2.531 (lowest)	2.563	2.554	2.544		



1 Calculate the frequency – used tally sheet.

weigth	tabulation	frequency	weigth	tabulation	frequency	weigth	tabulation	frequency
2.531		1	2.546	IIII	4	2.561	-	5
2.532	II	2	2.547		3	2.562		2
2.533		1	2.548		0	2.563		1
2.534		2	2.549		2	2.564		3
2.535		0	2.550		4	2.565		2
2.536		2	2.551	 	8	2.566		1
2.537		2	2.552	 	6	2.567		2
2.538		3	2.553		3	2.568	II	2
2.539		1	2.554		2	2.569		3
2.540		0	2.555		1	2.570		3
2.541		0	2.556	##	5	2.571	II	2
2.542		3	2.557		1	2.572		2
2.543	III	3	2.558	III	3	2.573		0
2.544		2	2.559	 	5	2.574		2
2.545	IIII	4	2.560	 	5	2.575		2



(2) Calculate the range.

$$R = X_h - X_l$$
= 2.575 - 2.531
= 0.044

3 Calculate the cell interval – use Sturgis' rule.

$$i = \frac{R}{1+3.322 \log n}$$

$$= \frac{0.044}{1+3.322 \log 110} = \frac{0.044}{1+3.322(2.041)} = 0.0057$$



- Trial and error
- Based on the guidelines, 0.005 will give the best presentation of the data.

$$i = 0.003;$$
 $h = \frac{R}{i} = \frac{0.044}{0.003} = 15$

$$i = 0.005;$$
 $h = \frac{R}{i} = \frac{0.044}{0.005} = 9$

$$i = 0.007;$$
 $h = \frac{R}{i} = \frac{0.044}{0.007} = 6$



4 Calculate the cell midpoint.


- The simplest technique is to select the lowest data point (2.531) as the midpoint value for the first cell.
- A better technique is to use the formula:

$$MP_l = X_l + \frac{i}{2} = 2.531 + \frac{0.005}{2} = 2.533$$



(5) Determine the cell boundaries.

- The boundary values are an extra decimal place or significant figure in accuracy than the observed values.
- Example:






6 Post the cell frequency.

Cell boundaries	Cell midpoint	frequency
2.5305 – 2.5355	2.533	6
2.5355 – 2.5405	2.538	8
2.5405 – 2.5455	2.543	12
2.5455 – 2.5505	2.548	13
2.5505 – 2.5555	2.553	20
2.5555 – 2.5605	2.558	19
2.5605 – 2.5655	2.563	13
2.5655 – 2.5705	2.568	11
2.5705 – 2.5755	2.573	8



7 Draw the histogram.






Frequency Polygon

- Relative frequencies of class intervals can also be shown in a frequency polygon.
- In a frequency distribution, the mid-value of each class is obtained.
- The frequency is plotted against the corresponding mid-value.
- These points are joined by straight lines.
- These straight lines may be extended in both directions to meet the X - axis to form a polygon.




Histogram & Polygon





Marks	Frequency
0 - 10	2
10 - 20	8
20 - 30	10
30 - 40	15
40 - 50	24
50 - 60	36
60 - 70	14
70 - 80	6
80 - 90	5




Polygon



Ogive

- A plot of the cumulative frequency against the upper class boundary with the points joined by line segments.
- Example:

				(Ogive)	(Relative Ogive)
Class num	Class	Frequency	Relative Frequency	Cumulative Freq.	Cumulative Relative Freq.
1	154 and under 161	4	0.15	4	0.15
2	161 and under 168	3	0.11	7	0.26
3	168 and under 175	5	0.19	12	0.44
4	175 and under 182	8	0.30	20	0.74
5	182 and under 189	6	0.22	26	0.96
6	189 and under 196	1	0.04	27	1.00
		27	1.00		





Stem-and-Leaf

- A stem-and-leaf display is an effective and compact way to summarize univariate numerical data.
- Each number in the data set is broken into two pieces, a stem and a leaf.
- The **stem** is the first part of the number and consists of the beginning digit(s).
- The **leaf** is the last part of the number and consists of the final digit(s).



When to use:

 Numerical data sets with a small to moderate number of observations (does not work well for very large data sets)

How to construct:

- Select one or more leading digits for the stem values. The trailing digits (or sometimes just the first one of the trailing digits) become the leaves.
- List possible stem values in a vertical column.
- Record the leaf for every observation beside the corresponding stem value.
- Indicate the units for stems and leaves someplace in the display.



What to look for:

The display conveys information about


- a representative or typical value in the data set
- the extent of spread about a typical value
- the presence of any gaps in the data
- the extent of symmetry in the distribution of values
- the number and location of peaks



Dataset:

63, 70, 74, 63, 82, 72, 95, 101, 86

High Temperature in Oswego (in °F)



Key: 6 / 3 means 63 °F

Stem: tens; leaf: ones



Exercise #3

a) Make a stem-and-leaf plot for the following list of numbers:

2.1, 2.0, 3.2, 3.7, 1.4, 1.5, 2.6

b) Make a stem-and-leaf plot for the following list of numbers:

23, 19, 28, 45, 36, 47, 48, 50, 51, 13



Box Plots

- Box plots a method that can summarize data that gives more detail than just a measure of center and spread and yet less detail than a stem-and-leaf display or histogram.
- It is compact, yet it provides information about center, spread, and symmetry or skewness of the data.




- Box plots graphically display five key statistics of a data set:
 - Minimum
 - First quartile
 - Median
 - Third quartile
 - Maximum
- Very useful in identifying the shape of a distribution and outliers in data set.



Quartiles

Quartiles are the values that divide a list of numbers into quarters.

- •First put the list of numbers in order.
- ■Then cut the list into four equal parts.
- ■The Quartiles are at the "cuts".





How To Calculate Quartile

 Sort the data set so measurements are in order from lowest to highest,

Calculate,

$$i = \frac{P}{100}(N)$$
 {Note: Q1: P = 25; Q2: P = 50; Q3: P = 75}

- If i is not an integer, round up to the next highest integer k and use Y[k] as the quartile estimate.
- If i is an integer, use $(Y[i] + Y[i+1]) \div 2$ as the quartile estimate.



Given set of data:

12, 4, 6, 11, 9, 15, 20, 18, 25, 30.

Calculate 1st, 2nd and 3rd quartile.

Solution (i): 1st quartile (Q1)

Arrange the data in ascending order:

4 6 9 11 12 15 18 20 25 30

•N=10, P=25; $i=25x10 \div 100=2.5$, k=3

Y[3] = 9, .: Q1 = 9



Solution (ii): 2nd quartile (Q2)

Arrange the data in ascending order:

4 6 9 11 12 15 18 20 25 30

• N=10, P=50; $i=50x10 \div 100=5$, k=5

$$(Y[5] + Y[6]) \div 2 = (12+15) \div 2 = 13.5, : Q2 = 13.5$$


Solution (Iii): 3rd quartile (Q3)

- Arrange the data in ascending order:
- N=10, P=75; $i=75x10 \div 100=7.5$, k=8

$$Y[8] = 20$$
, .: $Q3 = 20$



The "Interquartile Range" is from Q1 to Q3:





- 2 types of boxplots:
 - Skeletal boxplot
 - Modified boxplot




Skeletal Box Plots

- Construction of a Skeletal Boxplot
 - Draw a vertical (or horizontal) measurement scale
 - Construct a rectangular box with a lower (or left) edge at the lower quartile and a upper (or right) edge at the upper quartile.
 - The box width is then equal to the interquartile range (iqr)
 iqr = upper quartile lower quartile
 - Draw a horizontal (or vertical) line segment inside the box at the location of the median.
 - Extend vertical (or horizontal) line segments, call whiskers, from each of the box to the smallest and largest observations in the data set.



Example: Skeletal Box plot





Modified Box Plots

- An observation is an outlier if it is more than 1.5(iqr) away from the nearest quartile (the nearest end of the box)
- An outlier is **extreme** if it is more than 3(iqr) from the nearest end of the box, and it is **mild** otherwise.
- A modified boxplot represent mild and extreme outliers, and the whiskers extend on each end to the most extreme observations that are not outliers.



Outliers

 For a modified box plot, the whiskers are the lines that extend from the left and right of the box to the adjacent values. The adjacent values are defined as the lowest and highest observations that are still inside the region defined by the following limits:

○ Lower Limit: Q1 – 1.5 × IQR

○ Upper Limit: Q3 + 1.5 × IQR


• In general, values that fall outside of the adjacent value region are deemed outliers.



- Construction of a Modified Boxplot:
 - Draw a vertical (or horizontal) measurement scale
 - Construct a rectangular box with a lower (or left) edge at the lower quartile and a upper (or right) edge at the upper quartile.
 - The box width is then equal to iqr.
 - Draw a horizontal (or vertical) line segment inside the box at the location of the median.
 - Determine if there are any mild or extreme outliers in the data set.
 - Draw whiskers that extend from each end of the box to the most extreme observation that is not an outlier.
 - Draw a + to mark the location if any mild outliers or extreme outliers in the data set.



Example: Modified Box plot





Exercise #4

a) Draw a box plot for the following set of data. Remember to order the data first, if necessary.

1, 0, 3, 2, 1, 1, 7, 8, 6, 6, 7, 7

b) Draw a box plot for the following set of data.


4.7, 3.8, 3.9, 3.9, 4.6, 4.5, 5






Microsoft Excel:

Statistical Tools















R Programming:













LINPACK

Sales Dashboard

Top Performers


 Current Month
 Current Year
 Country
 Filter 3

 [Dec
 ▼
 [2019
 ▼
 [All)
 ▼
 [All]
 ▼

 Performance Scope
 defines color coding
 Customer
 Filter 4

 Current Month vs Same Month, Previous Year
 ▼
 [All)
 ▼
 [All]
 ▼

Dec-19 vs Dec-18



Lintao SA | Sales Dashboard | Confidential Information | For internal use only