

CHAPTER 8

POLAR COORDINATES

- 8.1 Polar coordinates system
- 8.2 Relationship between Cartesian and Polar Coordinates
- 8.3 Forming polar equations from Cartesian equations and vice-versa
- 8.4 Sketching polar equations
 - Method 1: Table
 - Method 2: Test of symmetries


8.1 Polar Coordinates System

Definition:

The polar coordinates of point P is written as an ordered pair (r, θ) , that is $P(r, \theta)$ where

r - distance from origin to P

θ - angle from polar axis to the line OP



Note:


- (i) θ is positive in anticlockwise direction, and it is negative in clockwise direction.
- (ii) Polar coordinate of a point is not unique.
- (iii) A point $(-r, \theta)$ is in the opposite direction of point (r, θ) .

Example: Plot the following set of points in the same diagram:


(a) $(3, 225^\circ)$, $(1, 225^\circ)$, $(-3, 225^\circ)$

(b) $(2, \frac{\pi}{3})$, $(2, -\frac{\pi}{3})$, $(-2, \frac{\pi}{3})$


a)



b)



For every point $P(r, \theta)$ in $0 \leq \theta \leq 2\pi$, there exist 3 more coordinates that represent the point P .




Example:

Find all possible polar coordinates of the points whose polar coordinates are given as the following:

- (a) $P(1, 45^\circ)$ (b) $Q(2, -60^\circ)$ (c) $R(-1, 225^\circ)$

8.2 Relationship between Cartesian and Polar Coordinates



1) Polar \implies Cartesian

$$x = r \cos \theta \quad y = r \sin \theta$$

2) Cartesian \implies Polar

$$r = \sqrt{x^2 + y^2} \quad \tan \theta = \frac{y}{x}$$

Example: Find the Cartesian coordinates of the points whose polar coordinates are given as

(a) $\left(1, \frac{7\pi}{4}\right)$ (b) $\left(-4, \frac{2\pi}{3}\right)$ (c) $(2, -30^\circ)$

Example: Find all polar coordinates of the points whose rectangular coordinates are given as

- (a) $(11,5)$ (b) $(0,2)$ (c) $(-4,-4)$

8.3 Forming polar equations from Cartesian equations and vice-versa.

To change the equation in Cartesian coordinates to polar coordinates, and conversely, use equation

$$x = r \cos \theta \quad y = r \sin \theta \quad r = \sqrt{x^2 + y^2}$$

Example: Express the following rectangular equations in polar equations.

(a) $y = x^2$

(b) $x^2 + y^2 = 16$

(c) $xy = 1$

Example: Express the following polar equations in rectangular equations.

(a) $r = 2 \sin \theta$

(b) $r = \frac{3}{4 \cos \theta + 5 \sin \theta}$

(c) $r = 4 \cos \theta + 4 \sin \theta$

(d) $r = \tan \theta \sec \theta$

8.4 Graph Sketching of Polar Equations

There are two methods to sketch a graph of $r = f(\theta)$:

(1) Form a table for r and θ . ($0 \leq \theta \leq 2\pi$).

From the table, plot the (r, θ) points.

(2) Symmetry test of the polar equation.

The polar equations is symmetrical

(a) about the x -axis if $f(-\theta) = f(\theta)$

- consider θ in range $[0, 180^0]$ only

(b) about the y -axis if $f(\pi - \theta) = f(\theta)$

- consider θ in range $[0, 90^0]$ **and** $[270^0, 360^0]$

(c) at the origin if $f(\pi + \theta) = f(\theta)$

- consider θ in range $[0, 180^0]$ **or** $[180^0, 360^0]$

* if symmetry at all, consider θ in range $[0, 90^0]$ **only**.


Example 1: Sketch the graph of $r = 2 \sin \theta$

Solution: (Method 1)

Here is the complete table

θ	0	30	60	90	120	150	180	210
$r = 2\sin\theta$	0	1.0	1.732	2	1.732	1	0	-1.0

θ	240	270	300	330	360
$r = 2\sin\theta$	-1.732	-2	-1.732	-1	0




Method 2

Symmetrical test for $f(\theta) = 2 \sin \theta$

Symmetry	Symmetrical test
About x-axis	$f(-\theta) = f(\theta) ?$
About y-axis	$f(\pi - \theta) = f(\theta) ?$
About origin	$f(\pi + \theta) = f(\theta) ?$

Since r symmetry at y -axis, consider θ in range $[0, 90^\circ]$ **and** $[270^\circ, 360^\circ]$

θ	0	30	60	90	270	300	330	360
$r = 2\sin\theta$	0	1.0	1.732	2	-2	-1.732	-1	0




Example 2: Sketch the graph of $r = \frac{3}{2} - \cos \theta$

Symmetry	Symmetrical test
About x-axis	$f(-\theta) = f(\theta) ?$
About y-axis	$f(\pi - \theta) = f(\theta) ?$
About origin	$f(\pi + \theta) = f(\theta) ?$

Since r symmetry at x -axis, consider θ in range $[0, 180^0]$ only.

θ	0	30	60	90	120	150	180
$r = \frac{3}{2} - \cos \theta$							



Example 3: Sketch the graph of $r = 2 \sin^2 \theta$

Symmetry	Symmetrical test
About x-axis	$f(-\theta) = f(\theta) ?$
About y-axis	$f(\pi - \theta) = f(\theta) ?$
About origin	$f(\pi + \theta) = f(\theta) ?$

Since r symmetry at _____, consider θ in range _____ only.

θ							
$r = 2\sin^2 \theta$							

