

Basic Counting Principles

- Counting principle is all about choices we might make given many possibilities.
- It is used to find the number of possible outcomes.
- It provides a basis of computing probabilities of discrete events.

innovative • entrepreneurial • global

2

OUTM

Basic Counting Principles

Some sample of counting problems:

- <u>Problem 1</u> How many ways are there to seat n couples at a round table, such that each couple sits together?
- <u>Problem 2</u>- How many ways are there to express a positive integer n as a sum of positive integers?

innovative • entrepreneurial • globa

vww.utm.my

TIM

Basic Counting Principles

Problem 3 - There are three boxes containing books. The first box contains 15 mathematics books by different authors, the second box contains 12 chemistry books by different authors, and the third box contains 10 computer science books by different authors. A student wants to take a book from one of the three boxes. In how many ways can the student do this?

innovative • entrepreneurial • global

www.utm.my

OUTM

Basic Counting Principles

<u>Problem 4</u> - The department will award a free computer to either a CS student or a CS professor. How many different choices are there, if there are 530 students and 15 professors?

innovative • entrepreneurial • global

www.utm.my

OUTM

<u>Problem 5</u> - A scholarship is available, and the student to receive this scholarship must be chosen from the Mathematics, Computer Science, or the Engineering Department. How many different choices are there for this student if there are 38 qualified students from the Mathematics Department, 45 qualified students from the Computer Science Department and 27 qualified students from the Engineering Department?

innovative • entrepreneurial • global

www.utm.my

Basic Counting Principles

- There are a number of basic principles that we can use to solve such problems:
 - 1) Addition Principle
 - 2) Multiplication Principle

7

UTM

Addition Principle

- Suppose that tasks T_1 , T_2 ,... T_k can be done in n_1 , n_2 ,... n_k ways, respectively.
- If all these tasks are **independent** of each other, then the number of ways to do one of these tasks is $n_1 + n_2 + ... + n_k$

innovative - entrepreneurial - alabal

..........

OUTM

Basic Counting Principles

■ If a task can be done in n_1 ways and a second task in n_2 ways, and if these two tasks cannot be done at the same time, then there are $n_1 + n_2$ ways to do either task.

innovative • entrepreneurial • glob

ww.utm.my

OUTM

Example 1

We want to find the number of integers between 5 and 50 that end with 1 or 7.

Solution:

- Let T denote this task. We divide T into the following tasks.
 - > T₁: find all integers between 5 and 50 that end with 1
 - ➤ T₂: find all integers between 5 and 50 that end with 7.

innovative • entrepreneurial • global

www.utm.my

UTM

- T_I : find all integers between 5 and 50 that end with 1.
 - →11, 21, 31, 41 => 4 ways
- T₂: find all integers between 5 and 50 that end with 7.
 - →7, 17, 27, 37, 47 => 5 ways
 - $.: n_1 + n_2 = 4 + 5 = 9$ ways.

innovative • entrepreneurial • global

..........

TIM UNITED TEXALOG BALANGE

Example 2

We want to find the number of integers between 4 and 100 that end with 3 or 5.

Solution:

Let *T* denote this task. We divide *T* into the following tasks.

- > T₁: find all integers between 4 and 100 that end with 3
- ➤ T₂: find all integers between 4 and 100 that end with 5.

innovative • entrepreneurial • glob

www.utm.my

TO UTM

- T_1 : find all integers between 4 and 100 that end with 3.
 - => 13, 23, 33, 43, 53, 63, 73, 83, 93
 - => 9 integers
- T₂: find all integers between 4 and 100 that end with 5.
 - => 5, 15, 25, 35, 45, 55, 65, 75, 85, 95
 - => 10 integers

innovative - entrepreneurial - alaba

w.utm.m


OUTM


- Task T_1 can be done in 9 ways.
- Task T₂ can be done in 10 ways.
- The number of ways to do one of these tasks is,


9+10=19


• Task T can be completed in 19 ways.

innovative • entrepreneurial • global









Multiplication Principle

A task T can be completed in k successive steps.

Step 1 can be completed in n_1 different ways. **Step 2** can be completed in n_2 different ways. **Step k** can be completed in n_k different ways.

■ Then the task T can be completed in $n_1.n_2....n_k$ different ways.

the state of the s

www.utm.my

TO UTM

Suppose that a procedure can be broken down into two successive tasks. If there are n_1 ways to do the first task and n_2 ways to do the second task after the first task has been done, then there are n_1n_2 ways to do the procedure.


innovative • entrepreneurial • global

20

OUTM

Example 4

Morgan is a lead actor in a new movie. She needs to shoot a scene in the morning in studio A and an afternoon scene in studio C. She looks at the map and finds that there is no direct route from studio A to studio C. Studio B is located between studios A and C. Morgan's friends Brad and Jennifer are shooting a movie in studio B. There are three roads, say A_1 , A_2 , and A_3 , from studio A to studio B and four roads, say B_1 , B_2 , B_3 , and B_4 , from studio B to studio C. In how many ways can Morgan go from studio A to studio C and have lunch with Brad and Jennifer at Studio B?



21 ww.utm.my

OUTM

• There are 3 ways to go from studio A to studio B and 4 ways to go from studio B to studio C.

≻T1 = 3

≻T2 = 4

 The number of ways to go from studio A to studio C via studio B is 3 × 4 = 12.

innovative • entrepreneurial • global

www.utm.my

OUTM

Example 5

- The letters *A*, *B*, *C*, *D*, and *E* are to be used to form strings of length 4.
- How many strings can be formed if we do not allow repetitions?
- For example:
 - BADE, ACBD, AEBC ..

innovative • entrepreneurial • global

www.utm.my


OUTM


- T_1 : choose the first letter \rightarrow 5 ways
- T_2 : choose the second letter \rightarrow 4 ways
- T_3 : choose the third letter \rightarrow 3 ways
- T_4 : choose the fourth letter \rightarrow 2 ways
- There are 5.4.3.2= 120 strings.


innovative • entrepreneurial • globa


www.utm.n


Example 6 The letters A, B, C, D, and E are to be used to form strings of length 4. How many strings can be formed if we allow repetitions? For example: — BABB, AABB, ACEE ...
























Basic Counting Principles
 The counting problem that we have considered so far involved either the addition principle or the multiplication principle.
 Sometimes, however, we need to use both of these counting principles to solve a particular problem.



Example

How many bit strings of length 8 either start with a 1 or end with 00?

Task 1: Construct a string of length 8 that starts with a

There is one way to pick the first bit (1), two ways to pick the second bit (0 or 1), two ways to pick the third bit (0 or 1),

. . two ways to pick the eighth bit (0 or 1).

•Product rule: Task 1 can be done in $1.2^7 = 128$ ways.

innovative • entrepreneurial • globa

TO UTM

Example (cont.)

Task 2: Construct a string of length 8 that ends with 00. There are two ways to pick the first bit (0 or 1), two ways to pick the second bit (0 or 1),

:

two ways to pick the sixth bit (0 or 1), one way to pick the seventh bit (0), and one way to pick the eighth bit (0).

Product rule: Task 2 can be done in $2^6 = 64$ ways.

annustina - antroneonousial - alabal

OUTM

Example (cont.)

Since there are 128 ways to do Task 1 and 64 ways to do Task 2, does this mean that there are 192 bit strings either starting with 1 or ending with 00?

No, because here Task 1 and Task 2 can be done at the same time.

When we carry out Task 1 and create strings starting with 1, some of these strings end with 00.

Therefore, we sometimes do Tasks 1 and 2 at the same time, so the sum rule does not apply.

innovative • entrepreneurial • global

www.utm.my

TIME UNITED TENNIOR BILLION

Example (cont)

If we want to use the sum rule in such a case, we have to subtract the cases when Tasks 1 and 2 are done at the same time.

How many cases are there, that is, how many strings start with 1 and end with 00?

There is one way to pick the first bit (1), two ways for the second, ..., sixth bit (0 or 1), one way for the seventh, eighth bit (0).

Product rule: In 2⁵ = 32 cases, Tasks 1 and 2 are carried out at the same time.

innovative • entrepreneurial • global

www.utm.my

OUTM

Example (cont)

Since there are 128 ways to complete Task 1 and 64 ways to complete Task 2, and in 32 of these cases Tasks 1 and 2 are completed at the same time, there are

128 + 64 - 32 = 160 ways to do either task.

innovative • entrepreneurial • global

www.utm.my


OUTM


Example 9


- The following items are available for breakfast.
 - ▶4 types of cereal
 - ▶2 types of juice
 - ➤3 types of bread
- How many ways a breakfast can be prepared if exactly 2 items are selected from 2 different groups?


innovative • entrepreneurial • global

www.utm.my










• In how many ways can this be done if either Aina or Wan must be chairperson?

➤ If Aina is chairperson 5. 4 = 20

➤ If Wan is chairperson 5. 4 = 20

➤ 20+20 = 40



TO UTM

- In how many ways can this be done if Syed must hold one of the position?
 - ➤ If Syed is chairperson 5. 4 = 20
 - ➤ If Syed is secretary 5. 4 = 20
 - ➤ If Syed is treasurer 5. 4 = 20
 - **>** 20+20+20 =60

innovative • entrepreneurial • globa

ww.utm.n

OUTM

- OR
 - ➤ Assign Syed for any position is 3 ways
 - Fill the highest remaining position is 5 ways
 - ➤ Fill the last position is 4 ways
 - **>**3. 5. 4 =60

innovative • entrepreneurial • global

www.utm.n

UTM

- In how many ways can this be done if Tan and Helmi must hold any position?
 - ➤ Assign Tan 3 ways
 - ➤ Assign Helmi 2 ways
 - Fill the remaining position 4 ways
 - **>**3. 2. 4 = 24

innovative • entrepreneurial • globa

ww.utm.my

TO UTM

Exercise

- How many eight-bit strings have either the second or the fourth bit 1(or both)?
- How many license plates of 2 letters from A to Z, followed by 3 digits from 0 to 9 can be made, if repetition of letters is not allowed?

innovative • entrepreneurial • global

OUTM

Exercise

 How many 5-digit telephone numbers can be constructed using digit 0-9 if each number starts with 67 and no digit appears more than once?

innovative • entrepreneurial • global

ww.utm.my