

UTM  
UNIVERSITI TEKNIK MALAYSIA

**SCSI1013: Discrete Structures**

**CHAPTER 1**

**Part 3:**  
**Fundamental and Elements of Logic**

2017/2018 – SEM. 1 : nzah@utm.my

innovative • entrepreneurial • global

1

www.utm.my

UTM  
UNIVERSITI TEKNIK MALAYSIA

**Why Are We Studying Logic?**

Some of the reasons:

- Logic is the foundation for computer operation
- Logical conditions are common in programs:  
**Example:**  
**Selection:** if (score <= max) { ... }  
**Iteration:** while (i < limit && list[i] != sentinel) ...
- All manner of structures in computing have properties that need to be proven (and proofs that need to be understood).  
**Examples:** Trees, Graphs, Recursive Algorithms, . . .
- Programs can be proven correct.
- Computational linguistics must represent and reason about human language, and language represents thought (and thus also logic).

innovative • entrepreneurial • global

2

www.utm.my

UTM  
UNIVERSITI TEKNIK MALAYSIA

**PROPOSITION**

A **statement** or a **proposition**, is a declarative sentence that is **either TRUE or FALSE, but not both**.

**Example:**

- 4 is less than 3.
- 7 is an even integer.
- Washington, DC, is the capital of United State.

innovative • entrepreneurial • global

3

www.utm.my

UTM  
UNIVERSITI TEKNIK MALAYSIA

**Example**

i) Why do we study mathematics?  
 ii) Study logic.  
 iii) What is your name?  
 iv) Quiet, please.

**The above sentences are not propositions. Why ?**

(i) & (iii) : is question, not a statement.  
 (ii) & (iv) : is a command.

innovative • entrepreneurial • global

4

www.utm.my

UTM  
UNIVERSITI TEKNIK MALAYSIA

**Example**

i) The temperature on the surface of the planet Venus is 800 F.  
 ii) The sun will come out tomorrow.

**Propositions? Why?**

i) Is a statement since it is either true or false, but not both.  
 ii) However, we do not know at this time to determine whether it is true or false.

innovative • entrepreneurial • global

5


www.utm.my

UTM  
UNIVERSITI TEKNIK MALAYSIA

**CONJUNCTIONS**

**Conjunctions** are:

- Compound propositions formed in English with the word “**and**”,
- Formed in logic with the caret symbol (“**∧**”), and
- True only when both participating propositions are true.



innovative • entrepreneurial • global

6

www.utm.my

**UTM** **CONJUNCTIONS** (cont.)

**TRUTH TABLE:** This tables aid in the evaluation of compound propositions.

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

innovative • entrepreneurial • global 7 www.utm.my

**UTM** **Example**

$p$  : 2 is an even integer } propositions  
 $q$  : 3 is an odd number }

$p \wedge q$  } symbols } statements  
 : 2 is an even integer and 3 is an odd number

$p$  : today is Monday ;  $q$  : it is hot

$p \wedge q$  : today is Monday and it is hot

innovative • entrepreneurial • global 8 www.utm.my

**UTM** **Example**

**Proposition:-**  
 $p$  : 2 divides 4  
 $q$  : 2 divides 6

**Symbol: Statement:-**  
 $p \wedge q$  : 2 divides 4 and 2 divides 6.  
 or,  
 $p \wedge q$  : 2 divides both 4 and 6.

innovative • entrepreneurial • global 9 www.utm.my

**UTM** **Example**


**Proposition:-**  
 $p$  : 5 is an integer  
 $q$  : 5 is not an odd integer

**Symbol: Statement:-**  
 $p \wedge q$  : 5 is an integer and 5 is not an odd integer.  
 or,  
 $p \wedge q$  : 5 is an integer but 5 is not an odd integer.

innovative • entrepreneurial • global 10 www.utm.my

**UTM** **DISJUNCTION**

- Compound propositions formed in English with the word "or",
- Formed in logic with the caret symbol (" $\vee$ "), and,
- True when one or both participating propositions are true.



innovative • entrepreneurial • global 11 www.utm.my

**UTM** **DISJUNCTION** (cont.)

Let  $p$  and  $q$  be propositions. The **disjunction** of  $p$  and  $q$ , written  $p \vee q$  is the statement formed by putting statements  $p$  and  $q$  together using the word "or". The symbol  $\vee$  is called "or"

innovative • entrepreneurial • global 12 www.utm.my

**UTM** **DISJUNCTION** (cont.)

The truth table for  $p \vee q$ :

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

innovative • entrepreneurial • global 13 www.utm.my

**UTM** **Example**

i)  $p$ : 2 is an integer ;  $q$ : 3 is greater than 5  
 $p \vee q \rightarrow$  2 is an integer or 3 is greater than 5

ii)  $p$ :  $1+1=3$  ;  $q$ : A decade is 10 years  
 $p \vee q \rightarrow$   $1+1=3$  or a decade is 10 years

iii)  $p$ : 3 is an even integer ;  $q$ : 3 is an odd integer  
 $p \vee q \rightarrow$  3 is an even integer or 3 is an odd integer ; or  
 3 is an even integer or an odd integer

innovative • entrepreneurial • global 14 www.utm.my

**UTM** **NEGATION**

Negating a proposition simply flips its value. Symbols representing negation include:  $\neg x, \bar{x}, \sim x, x'$  (NOT)

Let  $p$  be a proposition.  
 The negation of  $p$ , written  $\neg p$  is the statement obtained by negating statement  $p$ .

innovative • entrepreneurial • global 15 www.utm.my

**UTM** **NEGATION** (cont.)

The truth table of  $\neg p$ :

$p$	$\neg p$
T	F
F	T

innovative • entrepreneurial • global 16 www.utm.my

**UTM** **Example**

$p$ : 2 is positive  
 $\neg p \rightarrow$  2 is not positive.

innovative • entrepreneurial • global 17 www.utm.my

**UTM** **Exercise (1)**

$p$ : It will rain tomorrow ;  $q$ : it will snow tomorrow

Give the negation of the following statement and write the symbol.

It will rain tomorrow or it will snow tomorrow.

innovative • entrepreneurial • global 18 www.utm.my

**Exercise (2)**

In each of the following, form the conjunction and the disjunction of  $p$  and  $q$  by writing the symbol and the statements.

i)  $p$ : I will drive my car  
 $q$ : I will be late

ii)  $p$ :  $NUM > 10$   
 $q$ :  $NUM \leq 15$

**Exercise (3)**

Suppose  $x$  is a particular real number. Let  $p$ ,  $q$  and  $r$  symbolize " $0 < x$ ", " $x < 3$ " and " $x = 3$ ", respectively. Write the following inequalities symbolically:

a)  $x \leq 3$   
 b)  $0 < x < 3$   
 c)  $0 < x \leq 3$

**Solution:**  
 a)  $q \vee r$   
 b)  $p \wedge q$   
 c)  $p \wedge (q \vee r)$

**Exercise (4)**

State either TRUE or FALSE if  $p$  and  $r$  are TRUE and  $q$  is FALSE.

a)  $\sim p \wedge (q \vee r)$

a)  $(r \wedge \sim q) \vee (p \vee r)$

**CONDITIONAL PROPOSITIONS**

Let  $p$  and  $q$  be propositions.

**"if  $p$ , then  $q$ "**

is a statement called a **conditional proposition**, written as

$p \rightarrow q$

**CONDITIONAL PROPOSITIONS (cont.)**

The **truth table** of  $p \rightarrow q$   
 $\Rightarrow$  Cause and effect relationship

FALSE if  $p$  = True and  $q$  = false

$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

TRUE if both true or  $p$ =false for any value of  $q$

**Example**

$p$ : today is Sunday ;  $q$ : I will go for a walk

$p \rightarrow q$ : If today is Sunday, then I will go for a walk.

$p$ : I get a bonus ;  $q$ : I will buy a new car

$p \rightarrow q$ : If I get a bonus, then I will buy a new car

**Example**

$p$  :  $x/2$  is an integer.  
 $q$  :  $x$  is an even integer.

$p \rightarrow q$  : if  $x/2$  is an integer, then  $x$  is an even integer.

**BICONDITIONAL**

Let  $p$  and  $q$  be propositions.

**“ $p$  if and only if  $q$ ”**

is a statement called a **biconditional proposition**, written as

$p \leftrightarrow q$

**BICONDITIONAL (cont.)**

The **truth table** of  $p \leftrightarrow q$ :

$p$	$q$	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

**Example**

$p$  : my program will compile  
 $q$  : it has no syntax error.

$p \leftrightarrow q$  : My program will compile if and only if it has no syntax error.

$p$  :  $x$  is divisible by 3  
 $q$  :  $x$  is divisible by 9

$p \leftrightarrow q$  :  $x$  is divisible by 3 if and only if  $x$  is divisible by 9.

**LOGICAL EQUIVALENCE**

- The compound propositions  $Q$  and  $R$  are made up of the propositions  $p_1, \dots, p_n$ .
- $Q$  and  $R$  are logically equivalent and write,  $Q \equiv R$  provided that given any truth values of  $p_1, \dots, p_n$ , either  $Q$  and  $R$  are **both true** or  $Q$  and  $R$  are **both false**.

**Example**

$Q = p \rightarrow q$   
 $R = \neg q \rightarrow \neg p$   
 Show that,  $Q \equiv R$

The **truth table** shows that,  $Q \equiv R$

$p$	$q$	$p \rightarrow q$	$\neg q \rightarrow \neg p$
T	T	T	T
T	F	F	F
F	T	T	T
F	F	T	T

**Example**

Show that,  
 $\neg(p \rightarrow q) \equiv p \wedge \neg q$

The truth table shows that,  
 $\neg(p \rightarrow q) \equiv p \wedge \neg q$

$p$	$q$	$\neg(p \rightarrow q)$	$p \wedge \neg q$
T	T	F	F
T	F	T	T
F	T	F	F
F	F	F	F

**PRECEDENCE OF LOGICAL CONNECTIVES**

Precedence of logical connectives is as follows:

Connective	Symbol	Precedence
not	$\neg$	Highest
and	$\wedge$	
or	$\vee$	
if...then	$\rightarrow$	
If and only if	$\leftrightarrow$	Lowest

**Example**

Construct the truth table for,  
 $A = \neg(p \vee q) \rightarrow (q \wedge p)$

**Solution:**

$p$	$q$	$(p \vee q)$	$\neg(p \vee q)$	$(q \wedge p)$	$A$
T	T	T	F	T	T
T	F	T	F	F	T
F	T	T	F	F	T
F	F	F	T	F	F

**Exercise (5)**

Construct the truth table for each of the following statements:

- $\neg p \wedge q$
- $\neg(p \vee q) \rightarrow q$
- $\neg(\neg p \wedge q) \vee q$
- $(p \rightarrow q) \rightarrow (\neg q \rightarrow \neg p)$

**LOGIC & SET THEORY**

Logic and set theory go very well together. The previous definitions can be made very succinct:

- $x \notin A$  if and only if  $\neg(x \in A)$
- $A \subseteq B$  if and only if  $(x \in A \rightarrow x \in B)$  is True
- $x \in (A \cap B)$  if and only if  $(x \in A \wedge x \in B)$
- $x \in (A \cup B)$  if and only if  $(x \in A \vee x \in B)$
- $x \in A - B$  if and only if  $(x \in A \wedge x \notin B)$
- $x \in A \Delta B$  if and only if  $(x \in A \wedge x \notin B) \vee (x \in B \wedge x \notin A)$
- $x \in A'$  if and only if  $\neg(x \in A)$
- $X \in P(A)$  if and only if  $X \subseteq A$

**Venn Diagrams**

Venn Diagrams are used to depict the various unions, subsets, complements, intersections etc. of sets.

### Logic and Sets are closely related

**Tautology**

$p \vee q \leftrightarrow q \vee p$   
 $p \wedge q \leftrightarrow q \wedge p$   
 $p \vee (q \wedge r) \leftrightarrow (p \vee q) \wedge r$   
 $p \wedge (q \vee r) \leftrightarrow (p \wedge q) \vee r$   
 $p \vee (q \wedge r) \leftrightarrow (p \vee q) \wedge (p \vee r)$   
 $p \wedge (q \vee r) \leftrightarrow (p \wedge q) \vee (p \wedge r)$   
 $p \wedge \neg q \leftrightarrow p \wedge (\neg p \wedge q)$   
 $p \wedge (\neg q \vee r) \leftrightarrow (p \wedge \neg q) \wedge (p \wedge r)$   
 $p \wedge (\neg q \wedge r) \leftrightarrow (p \wedge \neg q) \vee (p \wedge r)$   
 $p \wedge (q \wedge \neg r) \leftrightarrow (p \wedge q) \wedge (\neg p \wedge r)$   
 $p \vee (q \wedge \neg r) \leftrightarrow (p \vee q) \wedge (\neg r \wedge \neg p)$   
 $p \wedge \neg \vee (q \wedge \neg r) \leftrightarrow (p \wedge \neg q) \vee (p \wedge r)$

**Set Operation Identity**

$A \cup B = B \cup A$   
 $A \cap B = B \cap A$   
 $A \cup (B \cap C) = (A \cup B) \cap C$   
 $A \cap (B \cup C) = (A \cap B) \cup C$   
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$   
 $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$   
 $A - B = A - (A \cap B)$   
 $A - (B \cap C) = (A - B) \cup (A - C)$   
 $A - (B \cup C) = (A - B) \cap (A - C)$   
 $A \cap (B - C) = (A \cap B) - (A \cap C)$   
 $A \cup (B - C) = (A \cup B) - (C - A)$   
 $A - (B - C) = (A - B) \cup (A \cap C)$

The above identities serve as the basis for an "algebra of sets".

### Logic and Sets are closely related

**Tautology**

$p \wedge p \leftrightarrow p$   
 $p \vee p \leftrightarrow p$   
 $p \wedge (\neg q \wedge \neg q) \leftrightarrow p$   
 $p \vee (\neg q \wedge \neg q) \leftrightarrow p$

**Contradiction**

$p \wedge \neg p$   
 $p \wedge (q \wedge \neg q)$   
 $p \wedge \neg p$

**Set Operation Identity**

$A \cap A = A$   
 $A \cup A = A$   
 $A - \emptyset = A$   
 $A \cup \emptyset = A$

**Set Operation Identity**

$A - A = \emptyset$   
 $A \cap \emptyset = \emptyset$   
 $A - A = \emptyset$

The above identities serve as the basis for an "algebra of sets".

### Theorem for Logic

Let  $p, q$  and  $r$  be propositions.

**Idempotent laws:**

$p \wedge p \equiv p$   
 $p \vee p \equiv p$

**Truth table:**

$p$	$p \wedge p$	$p \vee p$
T	T	T
F	F	F

### Theorem for Logic (cont.)

**Double negation law:**

$\neg \neg p \equiv p$

**Commutative laws:**

$p \wedge q \equiv q \wedge p$   
 $p \vee q \equiv q \vee p$

### Theorem for Logic (cont.)

**Associative laws:**

$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$   
 $(p \vee q) \vee r \equiv p \vee (q \vee r)$

**Distributive laws:**

$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$   
 $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$

**Absorption laws:**

$p \wedge (p \vee q) \equiv p$   
 $p \vee (p \wedge q) \equiv p$

### Prove: Distributive Laws

$p$	$q$	$r$	$p \vee (q \wedge r)$	$(p \vee q) \wedge (p \vee r)$
T	T	T	T	T
T	T	F	T	T
T	F	T	T	T
T	F	F	T	T
F	T	T	T	T
F	T	F	F	F
F	F	T	F	F
F	F	F	F	F

Prove: Absorption Laws

$p$	$q$	$p \wedge (p \vee q)$	$p \vee (p \wedge q)$
T	T	T	T
T	F	T	T
F	T	F	F
F	F	F	F

**Theorem for Logic** (cont.)

De Morgan's laws:

$$\neg(p \wedge q) \equiv (\neg p) \vee (\neg q)$$

$$\neg(p \vee q) \equiv (\neg p) \wedge (\neg q)$$

The truth table for  $\neg(p \vee q) \equiv (\neg p) \wedge (\neg q)$

$p$	$q$	$\neg(p \vee q)$	$\neg p \wedge \neg q$
T	T	F	F
T	F	F	F
F	T	F	F
F	F	T	T

**Exercise (6)**

Given,

$$R = p \wedge (\neg q \vee r)$$

$$Q = p \vee (q \wedge \neg r)$$

State whether or not  $R \equiv Q$ .

**Exercise (7)**

Propositional functions  $p$ ,  $q$  and  $r$  are defined as follows:

- $p$  is " $n = 7$ "
- $q$  is " $a > 5$ "
- $r$  is " $x = 0$ "

Write the following expressions in terms of  $p$ ,  $q$  and  $r$ , and show that each pair of expressions is **logically equivalent**. State carefully which of the above laws are used at each stage.

- $((n = 7) \vee (a > 5)) (x = 0)$   
 $((n = 7) (x = 0)) \vee ((a > 5) (x = 0))$
- $\neg((n = 7) (a \leq 5))$   
 $(n \neq 7) \vee (a > 5)$
- $(n = 7) \vee (\neg((a \leq 5) (x = 0)))$   
 $((n = 7) \vee (a > 5)) \vee (x \neq 0)$

**Exercise (8)**

Propositions  $p$ ,  $q$ ,  $r$  and  $s$  are defined as follows:

- $p$  is "I shall finish my Coursework Assignment"
- $q$  is "I shall work for forty hours this week"
- $r$  is "I shall pass Maths"
- $s$  is "I like Maths"

Write each sentence in symbols:

- I shall not finish my Coursework Assignment.
- I don't like Maths, but I shall finish my Coursework Assignment.
- If I finish my Coursework Assignment, I shall pass Maths.
- I shall pass Maths only if I work for forty hours this week and finish my Coursework Assignment.

Write each expression as a sensible (if untrue!) English sentence:

- $q \vee p$
- $\neg p \Rightarrow \neg r$

**Exercise (9)**

For each pair of expressions, construct **truth tables** to see if the two compound propositions are logically equivalent:

- $p \vee (q \wedge \neg p)$   
 $p \vee q$
- $(\neg p \wedge q) \vee (p \wedge \neg q)$   
 $(\neg p \wedge \neg q) \vee (p \wedge q)$