


SCSI2143: PROBABILITY & STATISTICAL DATA ANALYSIS

CHAPTER 3

Descriptive Statistics


Prepared & updated by: Razana/Suhaila/Nzah

innovative • entrepreneurial • global 2016/2017 – Semester 2 www.utm.my



Measures of Central Tendency


innovative • entrepreneurial • global 2 www.utm.my



Measurement of Central Tendency

- A measure of central tendency of a distribution is a numerical value that describes the central position of the data or how the data tend to build up in the center.
- Measurements:
 - Mean
 - Mode
 - Median

innovative • entrepreneurial • global 3 www.utm.my




Mean

- Mean is the sum of the observations divided by the number of observations.
- It is the most common measure of central tendency,

Sample mean: $\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$

Population mean: $\mu = \frac{\sum_{i=1}^N X_i}{N}$

innovative • entrepreneurial • global 4 www.utm.my




Example

- Data: 13, 18, 13, 14, 13, 16, 14, 21, 13

Calculation:
 $(13 + 18 + 13 + 14 + 13 + 16 + 14 + 21 + 13) \div 9 = 15$

- The mean is 15

innovative • entrepreneurial • global 5 www.utm.my



- The mean is unique for every set of data.
- Meaningful for interval and ratio data.
- Can be affected by outliers – rare observations that are radically different from the rest.

- Example: 3, 4, 6, 4, 7, 3, 6, 5, **1500**
- Mean: **170.89**

innovative • entrepreneurial • global 6 www.utm.my

Mean of Grouped Data

- The formula for the mean of grouped data,

$$\bar{X} = \frac{\sum_{i=1}^h f_i X_i}{n} = \frac{f_1 X_1 + f_2 X_2 + \dots + f_h X_h}{f_1 + f_2 + \dots + f_h}$$

where,

f_i : frequency in a class or frequency of an observed value.

X_i : class midpoint or an observed value.

n : number of classes or number of observed values.

Example

Find the mean value for the following data:

Number of children (X_i)	1	2	3	4	5	6	7
Frequency (f_i)	5	12	8	3	0	0	1

Solution:

$$\sum f_i X_i = 5(1) + 12(2) + 8(3) + 3(4) + 0(5) + 0(6) + 1(7) = 72$$

$$\bar{X} = \frac{\sum_{i=1}^h f_i X_i}{n} = \frac{72}{29} = 2.5$$

Example

Find the mean value for the following data:

Class interval	Frequency
40.5 – 45.5	7
45.5 – 50.5	10
50.5 – 55.5	15
55.5 – 60.5	2
60.5 – 65.5	6
Total	50

Example

Class interval	Midpoint	Frequency	$f_i X_i$
40.5 – 45.5	$(40.5 + 45.5) \div 2 = 43$	7	$43 \times 7 = 301$
45.5 – 50.5	48	10	480
50.5 – 55.5	53	15	795
55.5 – 60.5	58	2	696
60.5 – 65.5	63	6	378
Total		50	2650

$$\bar{X} = \frac{\sum_{i=1}^h f_i X_i}{n} = \frac{2650}{50} = 53$$

Median

- The median is the **middle** value when the data are arranged from smallest to largest.
- To find the median, your numbers have to be listed in an order, so you may have to rewrite your list first.
- For an odd number of observations, the formula for the place to find the median is

$$([\text{the number of data points}] + 1) \div 2$$

Example

- Data: 13, 18, 13, 14, 13, 16, 14, 21, 13
- Arrange in order: 13, 13, 13, 13, **14**, 14, 16, 18, 21
- There are nine numbers in the list, so the middle one will be the $(9 + 1) \div 2 = 10 \div 2 = 5^{\text{th}}$ number.
- So the median is 14.



- For an even number of observations, the median is the mean of the two middle numbers.
- Example:

2, 3, 3, **5**, **6**, 7, 8, 9

$$(5 + 6) \div 2 = \mathbf{5.5 \text{ (median)}}$$



- The median is meaningful for ratio, interval, and ordinal data.
- The median is not affected by outliers.

- Example: 3, 4, 6, 4, 7, 3, 6, 5, **1500**

3, 3, 4, 4, **5**, 6, 6, 7, 1500

- Median: **5**

Median of Grouped Data

- The median for the grouped data is given by

$$\text{median} = L + \frac{\frac{N}{2} - cf_p}{f_{med}}(W)$$

where,

(Median class is the first class with the value of cumulative frequency equal at least $N/2$)

L : lower class limit of median class,

N : total number of observations,

cf_p : cumulative frequency of the class preceding the median class,

f_{med} : frequency of the median class,

W : median class size.

Example

Class interval	Frequency	Cumulative frequency
40.5 – 45.5	7	7
45.5 – 50.5	10	17
50.5 – 55.5	15	32
55.5 – 60.5	2	34
60.5 – 65.5	6	40
Total	40	

$$N \div 2 = 40 \div 2 = 20$$

$$\therefore \text{median class} = 50.5 - 55.5$$

$$L = 50.5$$

$$N = 40$$

$$cf_p = 17$$

$$W = 5$$

$$f_{med} = 15$$

$$\text{median} = L + \frac{\frac{N}{2} - cf_p}{f_{med}}(W) = 51.5$$

Mode

- The mode is the value that occurs most often.
- If no number is repeated, then there is no mode for the list.

Example

Case study: On a cold winter day in January, the temperature for 9 North American cities is recorded in Fahrenheit as follows:

-8, 0, -3, 4, 12, 0, 5, -1, 0

What is the mode of these temperatures?

Solution:

- Ordering the data from least to greatest, we get:

-8, -3, -1, 0, 0, 0, 4, 5, 12

- The mode of these temperatures is 0.



Example

Case study: A marathon race was completed by 5 participants. The time taken by each participant is recorded as follows:

2.7 hr, 8.3 hr, 3.5 hr, 5.1 hr, 4.9 hr

What is the mode of these times given in hours?



Solution:

Ordering the data from least to greatest, we get:

2.7, 3.5, 4.9, 5.1, 8.3

Since each value occurs only once in the data set, there is **no mode** for this set of data.

Example

Case study: In a crash test, 11 cars were tested to determine what impact speed was required to obtain minimal bumper damage. The collected data as shown below:

24, 15, 18, 20, 18, 22, 24, 26, 18, 26, 24

Find the mode of the speeds given in miles per hour.

Solution:

Ordering the data from least to greatest, we get:



15, 18, 18, 18, 20, 22, 24, 24, 24, 26, 26

Since both 18 and 24 occur three times, the modes are 18 and 24 miles per hour. This data set is **bimodal**.

Mode of Grouped Data

- The first step towards finding the mode of the grouped data is to locate the class interval with the maximum frequency.
- The class interval corresponding to the maximum frequency is called the modal class.

- The mode of grouped data is calculated using the formula

$$\text{Mode} = l + h \times [(f_1 - f_0) \div (2f_1 - f_0 - f_2)]$$

where,


l : lower class limit of the modal class

h : class size assuming that all class intervals have the same class size

f_1 : frequency of the modal class

f_0 : frequency of the class preceding or just before the modal class

f_2 : frequency of the class succeeding or just after the modal class



Example

Number of Trees Planted (Class - Interval)	Number of Schools (Frequency: f_i)
5 - 25	12
25 - 45	8
45 - 65	14
65 - 85	20
85 - 105	6

The class interval corresponding to the maximum frequency is called the modal class.

$$\text{Mode} = l + h \times \frac{(f_1 - f_0)}{(2f_1 - f_0 - f_2)}$$


Where,


- l → lower class limit of the modal class.
- h → class size.
- f_1 → frequency of the modal class.
- f_0 → frequency of the class preceding or just before the modal class.
- f_2 → frequency of the class succeeding or just after the modal class.

In this case,

- $l = 65$
- $h = 20$
- $f_1 = 20$
- $f_0 = 14$
- $f_2 = 6$


innovative • entrepreneurial • global
23
www.utm.my





symmetrical distribution
asymmetrical distribution

innovative • entrepreneurial • global
24
www.utm.my



UTM
UNIVERSITI TEKNOLOGI MALAYSIA

Exercise #1

The owner of a shoe shop recorded the sizes of the feet of all the customers who bought shoes in his shop in one morning. These sizes are listed below:

8, 7, 4, 5, 9, 13, 10, 8, 8, 7, 6, 5, 3, 11, 10, 8, 5, 4, 8, 6

(a) What is the mean of these values?
 (b) What is the median of these values?
 (c) What is the mode of these values?

innovative • entrepreneurial • global

26

www.utm.my

Exercise #2

The table below gives the number of accidents each year at a particular road junction:

2001	2002	2003	2004	2005	2006	2007	2008
4	5	4	2	10	5	3	5

- Calculate the **mean**, **median** and **mode** for the values above.
- A road safety group want the council to do some improvement to make this junction safer. Which measure will they use to argue for this?
- The council don't want to spend money on the road junction. Which measure will they use to argue that safety work is not necessary?

Exercise #3

You grew fifty baby carrots using special soil. You dig them up and measure their lengths (to the nearest mm) and group the results:

Find the

- mean
- median
- mode

Length (mm)	Frequency
149.5 – 154.5	5
154.5 – 159.5	2
159.5 – 164.5	6
164.5 – 169.5	8
169.5 – 174.5	9
174.5 – 179.5	11
179.5 – 184.5	6
184.5 – 189.5	3

Exercise #4

Find mean, median and mode corresponding to the frequency table of samples of students cars and faculty/staff cars obtained from a college.

Age	Students	Faculty/Staff
0.5 – 3.5	23	30
3.5– 6.5	33	47
6.5 – 9.5	63	36
9.5 – 12.5	68	30
12.5 – 15.5	19	8
15.5 – 18.5	10	0
18.5 – 21.5	1	0
21.5 – 24.5	0	1


Data Profiles

- Percentile
- Quartile

Percentile

In a population or a sample, the P -th percentile is a value such that at least P percent of the values take on this value or less and at least $(100-P)$ percent of the values take on this value or more.

Example



- Sort the data set so measurements are in order from lowest to highest,

$$Y[1], Y[2], \dots, Y[N]$$

- Calculate,

$$i = \frac{P}{100}(N)$$

- If i is not an integer, round up to the next highest integer k and use $Y[k]$ as the percentile estimate.
- If i is an integer, use $(Y[i] + Y[i+1]) \div 2$ as the percentile estimate.

Example

Given set of data:

12, 4, 6, 11, 9, 15, 20, 18, 25, 30

- Calculate 80th percentile.
- Calculate 68th percentile.

Solution (i):

- Arrange in order:

4 6 9 11 12 15 18 20 25 30

- $N = 10, P = 80 ; i = 80 \times 10 \div 100 = 8$

$$Y[8] = 20, Y[9] = 25, P_{80} = (20 + 25) \div 2 = 22.5$$

Example

Solution (ii): Calculate 68th percentile

$N = 10, P = 68$

4 6 9 11 12 15 18 20 25 30

$i = 68 \times 10 \div 100 = 6.8, k = 7$

$Y[7] = 18, P_{68} = 18$

- The process of finding the percentile that corresponds to a particular value x is:

$$\text{percentile of value } x = \frac{\text{number of values less than } x}{\text{total number of values}} (100)$$

Example

Given a set of data as follows:

12 4 6 11 9 15 20 18 25 30

Find the percentile corresponding to the value,

$$Y[k] = 15$$

Solution: Arrange the data in order,



4 6 9 11 12 15 18 20 25 30


$$\text{percentile of value 15} = \frac{\text{number of values less than 15}}{\text{total number of values}}(100) = \frac{5}{10}(100) = 50$$

The value 15 is the 50th percentile.

Quartile

The 1st, 2nd, and 3d quartiles are the 25th, 50th, and 75th percentiles respectively.





Exercise #5

0.7901	0.8044	0.8062	0.8073	0.8079	0.8110
0.8126	0.8128	0.8143	0.8150	0.8150	0.8152
0.8152	0.8161	0.8161	0.8163	0.8165	0.8170


(a) Use the 18 sorted (left to right) weights of regular can drinks to find the percentile corresponding to the given value.

- i. 0.8143
- ii. 0.8062

(b) Find the indicated percentile and quartile.

- i. P_{80}
- ii. Q_3
- iii. P_{33}
- iv. Q_1


innovative • entrepreneurial • global 46 www.utm.my



UNIVERSITI TEKNOLOGI MALAYSIA

Measures of Dispersion & Shape

innovative • entrepreneurial • global 48 www.utm.my



UNIVERSITI TEKNOLOGI MALAYSIA

Measures of Dispersion

- Measures of dispersion measure how spread out a set of data is.
- **Range, variance, standard deviation**

innovative • entrepreneurial • global 49 www.utm.my


Range

- The range is the largest number in a set minus the smallest number.
- Example: 13, 18, 13, 14, 13, 16, 14, 21, 13

The largest value in the list is 21, and the smallest is 13,
so the range is $21 - 13 = 8$.


Variance

- A measure of the dispersion of a set of data points around their mean value.
- It is a mathematical expectation of the average squared deviations from the mean.



- Sample:
$$s^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n - 1}$$
- Population:
$$\sigma^2 = \frac{\sum_{i=1}^N (x_i - \mu)^2}{N}$$

innovative • entrepreneurial • global 52 www.utm.my



Standard Deviation

- A statistic used as a measure of the dispersion or variation in a distribution, equal to the square root of the arithmetic mean of the squares of the deviations from the arithmetic mean.
- It is the square root of the variance.

innovative • entrepreneurial • global 53 www.utm.my

Exercise #6

Bailey has been playing golf on the weekends for the past three years. Recently, she started keeping track of her recorded scores. Her scores for June and July at her favourite 9-hole (par 36) golf course are provided below:

**45 49 42 56 41 36 34 38 41 45 40 42 41 39
38 40 39 36 41**

Find the Range, Variance, and Standard Deviation for the above data.

Measures of Shape

- Skewness
- Kurtosis

Skewness

- Occurs when a distribution is not symmetrical about its mean.
- A distribution is symmetrical when its median, mean, and mode are equal.
- A positively skewed (**skewed to the right**) distribution occurs when the mean exceeds the median.
- A negatively skewed (**skewed to the left**) distribution occurs when the mean is less than the median.


Measuring Skewness

- Formula to measure skewness for univariate data x_1, x_2, \dots, x_N :


$$\text{Skewness} = \frac{\sum_{i=1}^N (x_i - \bar{x})^3}{(N-1)s^3}$$

Number of data points \rightarrow $(N-1)$
 s = std deviation \rightarrow s^3
 \bar{x} = mean \rightarrow $(x_i - \bar{x})^3$



- For normal distribution (symmetric distribution):
 - skewness = 0.
- Any symmetric data should have:
 - Skewness value near zero.
 - Distribution with mean, median and mode fall at the same point.






- Skewness > 0
 - The distribution is asymmetrical and points in the **positive direction**.
 - Example: Test scores of difficult examination where almost everyone did poorly on it
 - mode < median < mean



- Skewness < 0
 - The distribution is asymmetrical and points in the **negative direction**.
 - Example: Test scores of difficult examination where almost everyone did good on it
 - mode > median > mean







Kurtosis

- **Kurtosis** is the statistic which describes the degree of peakedness or flatness of a probability distribution relative to the benchmark normal distribution.
- In a similar way to the concept of skewness, *kurtosis* is a descriptor of the shape of a probability distribution
- Formula to measure kurtosis for univariate data x_1, x_2, \dots, x_N :

$$\text{Kurtosis} = \frac{\sum_{i=1}^N (x_i - \bar{x})^4}{(N-1)s^4}$$

UTM
UNIVERSITI TEKNOLOGI MALAYSIA

Excess Kurtosis

- Excess kurtosis is simply **kurtosis – 3**.
- A normal distribution has kurtosis exactly 3 (excess kurtosis exactly 0). Any distribution with kurtosis ≈ 3 (excess ≈ 0) is called **mesokurtic**.
- A distribution with kurtosis < 3 (excess kurtosis < 0) is called **platykurtic**. Compared to a normal distribution, its tails are shorter and thinner, and often its central peak is lower and broader.
- A distribution with kurtosis > 3 (excess kurtosis > 0) is called **leptokurtic**. Compared to a normal distribution, its tails are longer and fatter, and often its central peak is higher and sharper.

innovative • entrepreneurial • global 67 www.utm.my

