

UNIVERSITI TEKNOLOGI MALAYSIA

SCSI2143: PROBABILITY & STATISTICAL DATA ANALYSIS


CHAPTER 8

Analysis of Variance (ANOVA)

innovative • entrepreneurial • global

1

www.utm.my



UNIVERSITI TEKNOLOGI MALAYSIA

Outline

- Introduction
- One-Way ANOVA with Equal Sample Sizes.
- One-Way ANOVA with Unequal Sample Sizes.
- Two-Way ANOVA.

Not covered in this syllabus

innovative • entrepreneurial • global

2

www.utm.my



Introduction

- ANOVA is a method of testing the equality of three or more population means by analyzing sample variances.
- The purpose of ANOVA is to test for significant differences between Means.
- Elementary concepts provide a brief introduction to the basics of statistical significance testing.



One-Way ANOVA with Equal Sample Sizes

- We assume that the populations have normal distribution and same variance (or standard deviation), and the samples are random and independent of each other.


ANOVA Notation:

n = size of each sample

k = number of populations or treatments being compared

s_X^2 = variance sample means


s_p^2 = pooled variance obtained by calculating the mean of the sample variances



One-Way ANOVA with Equal Sample Sizes

- Test statistic:
$$F = \frac{\text{variance between sample}}{\text{variance within sample}} = \frac{nS_{\bar{X}}^2}{S_p^2}$$
- Variance between sample:
 - Also called variation due to treatment
 - An estimate of the common population variance σ^2 that is based on the variability among the sample **means**.
- Variance within sample:
 - Also called variation due to error.
 - An estimate of the common population variance σ^2 based on the sample **variances**.

innovative • entrepreneurial • global 5 www.utm.my



One-Way ANOVA with Equal Sample Sizes

- The critical value of F :
 - numerator degrees of freedom = $k - 1$
 - denominator degrees of freedom = $k(n - 1)$
 - k = number of population or treatments being compared.
 - n = sample size

innovative • entrepreneurial • global 6 www.utm.my

Example 1

Table below lists the head injury to car crash test dummies for four different types of cars. Use a 0.05 significance level to test the null hypothesis that the different types of car have the same mean.

No. of head injury			
Subcompact Cars	Compact Cars	Midsize Cars	Full-Size Cars
681	643	469	384
428	655	727	656
917	442	525	602
898	514	454	687
420	525	259	360


Example 1

- Step 1: Define hypothesis.

$$H_0 : \mu_1 = \mu_2 = \mu_3 = \mu_4$$

H_1 : at least one mean is different.

- Step 2: For each category, find n , \bar{x} and s



Example 1

- Category 1: (Subcompact Cars)

$$n = 5$$

$$\bar{x} = \frac{681 + 428 + 917 + 898 + 420}{5} = 668.8$$

$$s = \sqrt{\frac{(681 - 668.8)^2 + (428 - 668.8)^2 + (917 - 668.8)^2 + (898 - 668.8)^2 + (420 - 668.8)^2}{5 - 1}}$$

$$= 242.0$$
- Category 2: (Compact Cars)


$$n = 5$$

$$\bar{x} = \frac{643 + 655 + 442 + 514 + 525}{5} = 555.8$$

$$s = \sqrt{\frac{(643 - 555.8)^2 + (655 - 555.8)^2 + (442 - 555.8)^2 + (514 - 555.8)^2 + (525 - 555.8)^2}{5 - 1}}$$

$$= 91.0$$

innovative • entrepreneurial • global
www.utm.my



Example 1

- Category 3: (Midsize Cars)

$$n = 5$$

$$\bar{x} = \frac{469 + 727 + 525 + 454 + 259}{5} = 486.8$$

$$s = \sqrt{\frac{(469 - 486.8)^2 + (727 - 486.8)^2 + (525 - 486.8)^2 + (454 - 486.8)^2 + (259 - 486.8)^2}{5 - 1}}$$

$$= 167.7$$
- Category 4: (Fullsize Cars)

$$n = 5$$

$$\bar{x} = \frac{384 + 656 + 602 + 687 + 360}{5} = 537.8$$

$$s = \sqrt{\frac{(384 - 537.8)^2 + (656 - 537.8)^2 + (602 - 537.8)^2 + (687 - 537.8)^2 + (360 - 537.8)^2}{5 - 1}}$$

$$= 154.6$$

innovative • entrepreneurial • global
10
www.utm.my

Example 1

- Step 3: Find variance between samples

Step 3a: Find mean between samples

$$\bar{\bar{x}} = \frac{668.8 + 555.8 + 486.8 + 537.8}{k = 4} = 562.3$$

Step 3b: Find standard deviation between samples

$$s_{\bar{x}} = \sqrt{\frac{(668.8 - 562.3)^2 + (555.8 - 562.3)^2 + (486.8 - 562.3)^2 + (537.8 - 562.3)^2}{4 - 1}} \\ = 76.779$$

Step 3c: Find variance between samples

$$ns_{\bar{x}}^2 = 5(76.779)^2 = 29475.1$$

Example 1

- Step 4: Find variance within samples

$$s_p^2 = \frac{(242.0)^2 + (91.0)^2 + (167.7)^2 + (154.6)^2}{k = 4} \\ = 29717.4$$

- Step 5: Calculate test statistic, F

$$F = \frac{\text{variance between samples}}{\text{variance within samples}} = \frac{ns_{\bar{x}}^2}{s_p^2} \\ = \frac{29475.1}{29717.4} \\ = 0.992$$

Example 1

- Step 6: Calculate numerator and denominator degree of freedom
 - Numerator = $k - 1 = 4 - 1 = 3$
 - Denominator = $k(n - 1) = 4(5 - 1) = 16$
- Step 7: Find critical value of F with $\alpha = 0.05$ from F -distribution table
 - F critical value = 3.2389

Example 1

- Step 8: Test the claim and state the conclusion.
 - Since $F_{test\ statistic} < F_{critical\ value}$ ($0.992 < 3.2389$), we fail to reject the null hypothesis.
 - There is sufficient evidence to claim that the different types of cars have the same mean for head injury.

Example 2

Table below lists the chest deceleration to car crash test dummies for four different types of cars. Use a 0.05 significance level to test the null hypothesis that the different types of car have the same mean.

Subcompact Cars	Compact Cars	Midsize Cars	Full-Size Cars
n = 5	n = 5	n = 5	n = 5
$\bar{x} = 50.4$	$\bar{x} = 53.0$	$\bar{x} = 48.8$	$\bar{x} = 46.0$
s = 6.69	s = 4.64	s = 3.35	s = 7.11

Example 2

- Step 1: State the hypothesis.
 $H_0 : \mu_1 = \mu_2 = \mu_3 = \mu_4$
 $H_1 : \text{at least one mean is different.}$
- Step 2: (Done – refer table).
- Step 3: Find variance between samples

Step 3a: Find mean between samples

$$\bar{\bar{x}} = \frac{50.4 + 53.0 + 48.8 + 46.0}{4} = 49.55$$

Step 3b: Find standard deviation between samples

$$s_{\bar{x}} = \sqrt{\frac{(50.4 - 49.55)^2 + (53.0 - 49.55)^2 + (48.8 - 49.55)^2 + (46.0 - 49.55)^2}{4 - 1}}$$

$$= 2.93$$

Step 3c: Find variance between samples

$$ns_{\bar{x}}^2 = 5(2.93)^2 = 42.92$$

Example 2

- Step 4: Find variance within samples

$$s_p^2 = \frac{(6.69)^2 + (4.64)^2 + (3.35)^2 + (7.11)^2}{4}$$

$$= 32.02$$

- Step 5: Calculate test statistic, F

$$F = \frac{\text{variance between samples}}{\text{variance within samples}} = \frac{ns_{\bar{x}}^2}{s_p^2}$$

$$= \frac{42.92}{32.02}$$

$$= 1.34$$

Example 2

- Step 6: Calculate numerator and denominator degree of freedom
 - Numerator = $k - 1 = 4 - 1 = 3$
 - Denominator = $k(n - 1) = 4(5 - 1) = 16$
- Step 7: Find critical value of F with $\alpha = 0.05$ from F -distribution table
 - F critical value = 3.2389

Example 2

- Step 8: Test the claim and state the conclusion.
 - Since $F_{test\ statistic} < F_{critical\ value}$ ($1.34 < 3.2389$), we fail to reject the null hypothesis.
 - There is sufficient evidence to claim that the different types of cars have the same mean for chest deceleration.

Exercise

Table 1 lists the body temperatures of 5 randomly selected subjects from each of 3 different age groups. Informal examination of the 3 sample means (97.940, 98.580, 97.800) seems to suggest that the 3 samples come from populations with means that are not significantly different. Test the claim that the 3 age-group populations have the same mean body temperature. Use $\alpha = 0.05$.

Exercise

Table 1: Body Temperature (°F) Categorized by Age

18-20	21-29	30 and older
98.0	99.6	98.6
98.4	98.2	98.6
97.7	99.0	97.0
98.5	98.2	97.5
97.1	97.9	97.3

$n_1=5$	$n_2=5$	$n_3=5$
Mean: $\bar{x}_1=97.940$	$\bar{x}_2=98.580$	$\bar{x}_3=97.800$
Std. Dev: $s_1=0.568$	$s_2=0.701$	$s_3=0.752$