

## DISCRETE STRUCTURE (SECI 1013)

### TUTORIAL 1

**DUE DATE: 30 November, 2020**

- Let the universal set be the set  $\mathbf{R}$  of all real numbers and let  $A = \{x \in \mathbf{R} \mid 0 < x \leq 2\}$ ,  $B = \{x \in \mathbf{R} \mid 1 \leq x < 4\}$  and  $C = \{x \in \mathbf{R} \mid 3 \leq x < 9\}$ . Find each of the following:
  - $A \cup C$
  - $(A \cup B)'$
  - $A' \cup B'$
- Draw Venn diagrams to describe sets  $A$ ,  $B$ , and  $C$  that satisfy the given conditions.
  - $A \cap B = \emptyset$ ,  $A \subseteq C$ ,  $C \cap B \neq \emptyset$
  - $A \subseteq B$ ,  $C \subseteq B$ ,  $A \cap C \neq \emptyset$
  - $A \cap B \neq \emptyset$ ,  $B \cap C \neq \emptyset$ ,  $A \cap C = \emptyset$ ,  $A \not\subseteq B$ ,  $C \not\subseteq B$
- Given two relations  $S$  and  $T$  from  $A$  to  $B$ ,
$$S \cap T = \{(x, y) \in A \times B \mid (x, y) \in S \text{ and } (x, y) \in T\}$$
$$S \cup T = \{(x, y) \in A \times B \mid (x, y) \in S \text{ or } (x, y) \in T\}$$
Let  $A = \{-1, 1, 2, 4\}$  and  $B = \{1, 2\}$  and defined binary relations  $S$  and  $T$  from  $A$  to  $B$  as follows:

For all  $(x, y) \in A \times B$ ,  $x S y \leftrightarrow |x| = |y|$

For all  $(x, y) \in A \times B$ ,  $x T y \leftrightarrow x - y$  is even

State explicitly which ordered pairs are in  $A \times B$ ,  $S$ ,  $T$ ,  $S \cap T$ , and  $S \cup T$ .
- Show that  $\neg ((\neg p \wedge q) \vee (\neg p \wedge \neg q)) \vee (p \wedge q) \equiv p$ . State carefully which of the laws are used at each stage.
- $R_1 = \{(x, y) \mid x + y \leq 6\}$ ;  $R_1$  is from  $X$  to  $Y$ ;  $R_2 = \{(y, z) \mid y > z\}$ ;  $R_2$  is from  $Y$  to  $Z$ ; ordering of  $X$ ,  $Y$ , and  $Z$ : 1, 2, 3, 4, 5.

Find:

  - The matrix  $A_1$  of the relation  $R_1$  (relative to the given orderings)
  - The matrix  $A_2$  of the relation  $R_2$  (relative to the given orderings)
  - Is  $R_1$  reflexive, symmetric, transitive, and/or an equivalence relation?
  - Is  $R_2$  reflexive, antisymmetric, transitive, and/or a partial order relation?
- Suppose that the matrix of relation  $R_1$  on  $\{1, 2, 3\}$  is

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}$$

relative to the ordering 1, 2, 3, and that the matrix of relation  $R_2$  on  $\{1, 2, 3\}$  is

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

relative to the ordering 1, 2, 3. Find:

- a) The matrix of relation  $R_1 \cup R_2$
  - b) The matrix of relation  $R_1 \cap R_2$
7. If  $f: \mathbf{R} \rightarrow \mathbf{R}$  and  $g: \mathbf{R} \rightarrow \mathbf{R}$  are both one-to-one, is  $f + g$  also one-to-one? Justify your answer.
  8. With each step you take when climbing a staircase, you can move up either one stair or two stairs. As a result, you can climb the entire staircase taking one stair at a time, taking two at a time, or taking a combination of one- or two-stair increments. For each integer  $n \geq 1$ , if the staircase consists of  $n$  stairs, let  $c_n$  be the number of different ways to climb the staircase. Find a recurrence relation for  $c_1, c_2, \dots, c_n$ .
  9. The Tribonacci sequence  $(t_n)$  is defined by the equations,

$$t_1 = t_2 = t_3 = 1, \quad t_n = t_{n-1} + t_{n-2} + t_{n-3} \quad \text{for all } n \geq 4.$$

- a) Find  $t_7$ .
- b) Write a recursive algorithm to compute  $t_n, n \geq 1$ .